

REVIEW ARTICLE

Low Dose Radiation Therapy for Covid-19: Benefit or Threat?

Ahlam Almanie, Samaneh Abbasi

Ahlam Almanie

Abadan Faculty of Medical Sciences
Faculty of Medical Sciences, Abadan, Iran
Email: maalmanni90@gmail.com

Samaneh Abbasi

Faculty of Medical Sciences, Abadan, Iran.

ABSTRACT

The new coronavirus outbreak emerged at the end of 2019 worldwide in a very short period of time. The number of victims of this virus until 2020/11/17 has been 1323143. The consequences of these viruses in humans are common cold or mild illness in the upper respiratory region. In more severe cases can causes severe interstitial pneumonia and acute respiratory distress syndrome (ARDS) and Middle East respiratory syndrome (MERS) and, severe acute respiratory syndrome (SARS). So far various drugs have been prescribed and used for the treatment. However, their efficiency and their side effects for treatment of pneumonia of COVID19 are unknown and should be more investigated. Low-dose radiation (LDR) (30 to 100 cGray(Gy)) has been used historically since the early 1930s with hopeful results for pneumonia treatment and was a common treatment solution for viral pneumonia until the 1940s.

As some recent studies have raised the use of LDRT for COVID19 treatment, we sought to review previous evidence oftherapeutic role of LDRT in the inflammatory diseases as well as recent recommendations about consider LDRT as the treatment method for COVID19.

Based on the available evidence and the background of studies, it seems that choosing a dose 0.3-0.5 Gy in severe cases of the disease, as well as using radiation for the whole body instead of the lungs, can optimize the immune system, and optimizing the immune system will help improve COVID-19.

KEYWORDS: Coronavirus, LDRT, X-ray, Radiotherapy

INTRODUCTION

The new coronavirus outbreaks emerged at the end of 2019 to the entire world in a very short time. The number of victims of this virus until 2020/11/17 has been 1323143¹. The Coronavirus (CoV) is a single-stranded, RNA viruses that can infect animals and humans² and, like other RNA viruses are mutating³. At first, the new virus was named 2019-nCoV⁴, subsequently, the International Committee on Taxonomy of Viruses (ICTV) applied the term SARS-CoV-2 for the virus and COVID-19 for the disease⁴.

The consequences of these viruses can be in humans, common cold or mild illness in the upper respiratory region. In more severe cases can cause severe interstitial pneumonia and acute respiratory distress syndrome (ARDS) and the middle east respiratory syndrome (MERS) and, severe acute respiratory syndrome (SARS)^{5,6}. In acute cases, the virus cannot be controlled effectively by the immune system and can spread more widely, this lead to activate macrophages and granulocytes and release pro-inflammatory cytokines and lung tissue damage². Proinflammation and secretion of cytokines, including IL-6, occur in COVID-19 (7). To control inflammation leading to ARDS and respiratory failure, several promising medicines that directly or indirectly play an anti-inflammatory and inhibitory role in IL-6 have been investigated⁷⁻⁹.


The WHO has confirmed that currently there isn't any safe and effective drug¹⁰ but only supportive drugs¹¹. So far, various drugs have been prescribed and used for treatment. New research shows that antiviral drugs that inhibit neuraminidase (Oseltamivir, Peramivir Zanamivir, Ganciclovir, Aciclovir, Ribavirin) are ineffective against COVID19¹². There is some evidence about the effectiveness of the (Remdesivir, Lopinavir, Ritonavir, Lopinavir/ Ritonavir Combined with interferon- β , plasma therapy, and monoclonal antibodies for COVID19. However, their efficiency and their side effects for the treatment of pneumonia of COVID19 are unknown and should be more investigated¹².

Besides, it may not be necessary to take medicine at the early stages of the disease and may weaken the immune system and cause viral replication. The use of drugs in an advanced stage of the disease is still debatable¹³. Improper use of the drug may also lead to rapid adaptation to resistance¹⁴. The provision of medicine and treatment economically has put a lot of pressure on the treatment system of all countries. Therefore, it is necessary to find an effective treatment method that is both economically viable and reliable.

At the beginning of the discovery of X-rays, its diagnostic application was more considered, but later doctors noticed significantly changes in the patient's medical condition by radiation, and it is considered as a therapeutic applications¹⁵.

Low-dose radiation (LDR) (30 to 100 centiGray(cGy)) has been used historically since the early 1930s with hopeful results for pneumonia treatment and was a common treatment solution for viral pneumonia up to the 1940s¹⁶⁻¹⁸. The low dose of radiation (LDR) is a dose that creates multiple collisions and sufficient reactive oxygen species (ROS) to actuate the immune system and cause visible health advantages. High-dose radiation(HDR) is a dose that suppresses and damages the immune system, causing immediate or latent damage (Figure I)¹⁵.

FIGURE I: DEFINITION OF A LOW DOSE AND A HIGH DOSE OF RADIATION¹⁵



HDR Vs LDR

There is strong evidence, in contrast, high-dose radiation that produces inflammatory cytokines into immune and endothelial cells¹⁹ low-dose radiation, Induces anti-inflammatory effects through a various mechanism such as inducing apoptosis in immune cells, secreting anti-inflammatory factors and reducing macrophage function, and preventing cytokine storms^{13,20,21} and reduction of inflammatory cytokines such as IL-6²². Low-dose radiation therapy's (LDRT) anti-inflammatory effect reaches its maximum 48 hours after irradiation and disappears after 72 hours²²⁻²⁵.

There are also studies showing that LDRT can improve the immune response in animals²⁶. Animal models show LDRT can halve the acute phase of pneumonia¹⁸. Over the past decade, the modulation of many immunological procedures by LDRT has been studied in vitro and in vivo. The most effective modulation of these processes occurs at doses of 0.3-0.7 Gy.²³ LDRT (0.5 Gy per fraction) also reduces inflammation in musculoskeletal disorders²⁷.

Mechanism of action of LDRT

A review of a study on the therapeutic effects of LDRT on bacterial pneumonia (lobular and bronchopneumonia), sulfanilamide non-responsive, interstitial, and atypical pneumonia, especially when irradiation was performed on days 6-14, clinical symptoms started, answered appropriately and lead to rapidly relieving symptoms, increased recovery, and reduced mortality¹⁶.

Various environmental factors polarize macrophages into two separate phenotypes: a pro-oxidative/pro-inflammatory M1 phenotype and an anti-inflammatory M2 phenotype²⁸. Calabrese considers the polarization of macrophages to an anti-inflammatory or M2 phenotype a possible the process by which radiation therapy(RT) decreases inflammation and simplifies recovery²⁹.

LDR, on the other hand, can increase the toxicity of Natural Killer cells (NK cells) and differentiate mature dendritic cells (mDCs) and M1 macrophages and activate T helper type 1 (Th1) and B cells, which improve the immune system and LDR regulates the negative effects of the immune response by converting immature DCs to mDCs and inducing M2F macrophages differentiation and stimulating Regulatory T cells(Treg) retention/expansion³⁰.

A study on rats showed that LDRT regulates immune responses that exhibit immune hormone features. Multiple molecular functions and cellular components are involved in the clinical efficacy of LDRT³¹.

Past studies also show that LDRT regulates the number of lymphocytes, control bacterial infections, and can regulate excessively inflammatory responses, exactly what COVID-19 patients need³².

Use of LDRT in the treatment of Covid-19

Recently, a team of American and Iranian researchers proposed the use of LDRT to treat COVID-19. Kirkby and Mackenzie also said that pneumonia treatment with X-rays is not a new issue in the 20th century and treat the lungs affected by pneumonia caused by COVID-19 with low linear energy transfer(LET)and dose fewer than 100cGy radiation has been shown to reduce inflammation and relieve life-threatening symptoms³³.

Several medical institutions have begun radiation therapy for COVID-19 and several LDR tests to the lungs were being performed in the United States for people with COVID-19 pneumonia. And there are several similar cases internationally³⁴.

The countries undergoing these tests are the United States, Spain, Italy, Iran, France, Belgium, and India³⁵.

A clinical study is performing on COVID19 patients in Spain, with a dose of 0.5Gy to lung at a single fraction of 1 Gy in 2 separate fractions 48–72 h after the first. Judgment is based on measuring the PaFio2 ratio [the ratio of the partial pressure of oxygen, PaO2 and the percentage of oxygen supplied, FiO2) 48-72 hours after irradiation, comparing radiographs, molecular adhesion, anti-inflammatory cytokines, and oxidative stress mediators. The results of this study have not yet been published¹⁹.

It is important to note that in recent reports, in 20% -30% of patients, blood clotting has been reported³⁶⁻³⁸. Which oxidative can reduce or block blood clotting³⁹ So instead of irradiating the lungs, the whole body should be irradiated⁴⁰.

Concerns about the consequences

Despite all evidence that supports the anti-inflammatory effects of LDRT, but some scientist raised concerns about the suppression of immune response against infectious agents due to the anti-inflammatory effects of LDRT in the treatment of COVID-19. And also Since there is some evidence about the possibility of the spread and transcription of viruses due to the use of radiation therapy⁴¹⁻⁴⁴, they have suggested that the whole body be irradiated at the right time instead of only lung radiation⁴⁵.

In response to these concerns, Carles agrees with the need for preclinical and clinical examination of this procedure and the importance of the timegiving this dose. However, he believes that the risks posed by this study are very small and the induction of secondary diseases for doses less than 600cGy is negligible⁴⁶. Evidence suggests that the threshold dose that induces secondary cancer is higher than 1.1 Gy¹⁵.

Some scientists believe that LDR can induce anti-cancer effects and inhibit tumor growth and metastasis. They even believe DNA repair ability and T-cell reactive ability, in people who live in a high level of natural radiation(HNR) area are increased⁴⁷. Since computed tomography (CT) dose is 5cGy and the recommended dose for treatment is 6 to 20 CT, the risk of secondary complications is very small¹³.

The safety of this method has been investigated by various studies, which show the risk of radiation effects with low doses is very small⁴⁸⁻⁵¹. Despite all the previous studies on the treatment of non-malignant diseases⁵² and the induction of anti-inflammatory effects in animals

by LDRT³¹, it is not possible to determine the correct time of irradiation and the start and end time of treatment in COVID-19. Therefore, the need for preclinical studies is undeniable. There are also problems such as the workload pressure on radiotherapy devices and the problem of isolating cancer patients from COVID-19 patients³³.

CONCLUSION

LDRT is an affordable non-toxic treatment that is available in most public hospitals. So It can be used for a large number of patients who do not receive specific anti-IL-6 treatments at the intensive care units (ICU) in low and middle income countries. Based on the available evidence and the background of studies, it seems that choosing a dose 0.3-0.5 Gy in severe cases of the disease, as well as using radiation for the whole body instead of the lungs, can optimize the immune system, and optimizing the immune system will help improve COVID-19.

REFERENCES

1. South China Newspost. Coronavirus pandemic current news. Confirmed Covid-19 Cases. [Internet]. Available from <https://multimedia.scmp.com/widgets/china/wuhanvirus/> .
2. Conti P, Ronconi G, Caraffa A, Gallenga C, Ross R, Frydas I, et al. Induction of pro-inflammatory cytokines (IL-1 and IL-6) and lung inflammation by Coronavirus-19 (COVI-19 or SARS-CoV-2): anti-inflammatory strategies. *J Biol Regul Homeost Agents*. 2020; 34(2): 1.
3. Rödel F, Arenas M, Ott OJ, Fournier C, Georgakilas AG, Tapio S, et al. Low-dose radiation therapy for COVID-19 pneumopathy: what is the evidence? *Strahlentherapie und Onkologie*. 2020: 1-4.
4. Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R. COVID-19 infection: Origin, transmission, and characteristics of human coronaviruses. *J Adv Res*. 2020.
5. Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, evaluation and treatment coronavirus (COVID-19). *Statpearls [internet]: StatPearls Publishing*; 2020.
6. Petrosillo N, Viceconte G, Ergonul O, Ippolito G, Petersen E. COVID-19, SARS and MERS: are they closely related? *Clinical Microbiology and Infection*. 2020.
7. Russell B, Moss C, George G, Santaolalla A, Cope A, Papa S, et al. Associations between immune-suppressive and stimulating drugs and novel COVID-19—a systematic review of current evidence. *E Cancer Med Sci*. 2020; 14.
8. Mehta P, McAuley DF, Brown M, Sanchez E, Tattersall RS, Manson JJ, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet (London, England)*. 2020; 395(10229): 1033.
9. Siddiqi HK, Mehra MR. COVID-19 illness in native and immunosuppressed states: A clinical–therapeutic staging proposal. *J Heart Lung Transplant*. 2020; 39(5): 405.
10. Young K. NIH Publishes COVID-19 Treatment Guidelines. *J Watch*. 2020.
11. Deng S-Q, Peng H-J. Characteristics of and public health responses to the coronavirus disease 2019 outbreak in China. *J Clin Med*. 2020; 9(2): 575.
12. Li H, Wang Y, Xu J, Cao B. Potential antiviral therapeutics for 2019 Novel Coronavirus. *Chinese J Tuberculos Respirat Dis*. 2020; 43: E002-E.
13. Montero A, Arenas M, Algara M. Low-dose radiation therapy: could it be a game-changer for COVID-19? *Clin Translat Oncol*. 2020: 1.
14. Irwin KK, Renzette N, Kowalik TF, Jensen JD. Antiviral drug resistance as an adaptive process. *Virus evolution*. 2016; 2(1).
15. Cuttler JM. Application of low doses of ionizing radiation in medical therapies. *Dose-Response*. 2020; 18(1): 1559325819895739.
16. Calabrese EJ, Dhawan G. How radiotherapy was historically used to treat pneumonia: could it be useful today? *Yale J Biol Med*. 2013; 86(4): 555.
17. Oppenheimer A. Roentgen therapy of “virus” pneumonia. *Am J Roentgenol Radiat Ther*. 1943; 49: 635-8.
18. Dubin I, Baylin G, GOBBLE JW. The effect of roentgen therapy on experimental virus pneumonia; on pneumonia produced in white mice by swine influenza virus. *Am J Roentgenol Radium Ther*. 1946; 55: 478-81.
19. Algara M, Arenas M, Marin J, Vallverdu I, Fernandez-Letón P, Villar J, et al. Low dose anti-inflammatory radiotherapy for the treatment of pneumonia by covid-19: a proposal for a

- multi-centric prospective trial. *Clin Translat Radiat Oncol*. 2020.
20. Arenas M. Anti-inflammatory effects of low-dose radiotherapy: Indications, dose and radiobiological mechanisms involved. *Rep Pract Oncol Radiother*. 2013; 18: S12-S3.
 21. Rödel F, Keilholz L, Herrmann M, Sauer R, Hildebrandt G. Radiobiological mechanisms in inflammatory diseases of low-dose radiation therapy. *Int J Radiat Biol*. 2007; 83(6): 357-66.
 22. Deloch L, Fuchs J, Rückert M, Fietkau R, Frey B, Gaipl US. Low-dose irradiation differentially impacts macrophage phenotype in dependence of fibroblast-like synoviocytes and radiation dose. *J Immunology Res*. 2019; 2019.
 23. Rödel F, Frey B, Manda K, Hildebrandt G, Hehlhans S, Keilholz L, et al. Immunomodulatory properties and molecular effects in inflammatory diseases of low-dose x-irradiation. *Front Oncol*. 2012; 2: 120.
 24. Royo LT, Redondo GA, Pianetta MÁ, Prat MA. Low-dose radiation therapy for benign pathologies. *Reports of Practical Oncology & Radiotherapy*. 2020.
 25. Lödermann B, Wunderlich R, Frey S, Schorn C, Stangl S, Rödel F, et al. Low dose ionising radiation leads to a NF- κ B dependent decreased secretion of active IL-1 β by activated macrophages with a discontinuous dose-dependency. *Int J Radiat Biol*. 2012.
 26. Yang G, Kong Q, Wang G, Jin H, Zhou L, Yu D, et al. Low-dose ionizing radiation induces direct activation of natural killer cells and provides a novel approach for adoptive cellular immunotherapy. *Cancer Biotherapy and Radiopharmaceut*. 2014; 29(10): 428-34.
 27. Ott OJ, Niewald M, Weitmann H-D, Jacob I, Adamietz IA, Schaefer U, et al. DEGRO guidelines for the radiotherapy of non-malignant disorders. *Strahlentherapie und Onkologie*. 2015; 191(1): 1-6.
 28. Calabrese EJ, Giordano JJ, Kozumbo WJ, Leak RK, Bhatia TN. Hormesis mediates dose-sensitive shifts in macrophage activation patterns. *Pharmacol Res*. 2018; 137: 236-49.
 29. Calabrese E, Dhawan G, Kapoor R, Kozumbo W. Radiotherapy treatment of human inflammatory diseases and conditions: optimal dose. *Human & experimental toxicology*. 2019; 38(8): 888-98.
 30. Cui J, Yang G, Pan Z, Zhao Y, Liang X, Li W, et al. Hormetic response to low-dose radiation: Focus on the immune system and its clinical implications. *Int J Molecul Sci*. 2017; 18(2): 280.
 31. El-fatah A, Abd El-wahab H, Ezz MK, El-kabany HA, ElSonbaty SM. Reduction of some extra-articular complications associated with arthritis development in rats by low dose γ -irradiation. *Arab J Nuclear Sci Applic*. 2020; 53(1): 172-81.
 32. Ghadimi-Moghadam A, Haghani M, Bevelacqua J, Jafarzadeh A, Kaveh-Ahangar A, Mortazavi S, et al. COVID-19 tragic pandemic: concerns over unintentional “directed accelerated evolution” of novel Coronavirus (SARS-CoV-2) and introducing a modified treatment method for ARDS. *J Biomed Physics Engineer*. 2020; 10(2): 241.
 33. Kirkby C, Mackenzie M. Is low dose radiation therapy a potential treatment for COVID-19 pneumonia? *Radiotherapy and Oncology*. 2020.
 34. Conca J. Preliminary Data Suggests Low-Dose Radiation May Be Successful Treatment For Severe Covid-19 2020
 35. <https://clinicaltrials.gov/> [Internet].
 36. Willyard C. Coronavirus blood-clot mystery intensifies. *Nature*. 2020.
 37. Klok FA, Kruip M, Van Der Meer N, Arbous M, Gommers D, Kant K, et al. Confirmation of the high cumulative incidence of thrombotic complications in critically ill ICU patients with COVID-19: an updated analysis. *Thrombosis Res*. 2020.

38. Poissy J, Goutay J, Caplan M, Parmentier E, Duburcq T, Lassalle F, et al. Pulmonary embolism in COVID-19 patients: awareness of an increased prevalence. *Circulation*. 2020.
39. Violi F, Pignatelli P. Platelet oxidative stress and thrombosis. *Thrombosis Res*. 2012; 129(3): 378-81.
40. Mehdizadeh A, Bevelacqua J, Mortazavi S, Mortazavi S. COVID-19: Introducing Low Dose Radiation as an Effective Treatment for Pneumonia that Shouldn't Induce Selective Pressure and New Mutations. *J Biomed Phys Eng*. 2020; 10(3): 247-50.
41. Ceccaldi P. Ionizing radiation modulates the spread of an apathogenic rabies virus in mouse brain. *Int J Radiat Biol*. 1996; 70(1): 69-75.
42. Ramirez-Fort MK, Zeng J, Feily A, Ramirez-Pacheco LA, Jenrette JM, Mayhew DL, et al. Radiotherapy-induced reactivation of neurotrophic human herpes viruses: Overview and management. *J Clin Virol*. 2018; 98: 18-27.
43. Mezhir JJ, Advani SJ, Smith KD, Darga TE, Poon AP, Schmidt H, et al. Ionizing radiation activates late herpes simplex virus 1 promoters via the p38 pathway in tumors treated with oncolytic viruses. *Cancer Research*. 2005; 65(20): 9479-84.
44. Iordanskiy S, Van Duyne R, Sampey GC, Woodson CM, Fry K, Saifuddin M, et al. Therapeutic doses of irradiation activate viral transcription and induce apoptosis in HIV-1 infected cells. *Virology*. 2015; 485: 1-15.
45. Kefayat A, Ghahremani F. Low dose radiation therapy for COVID-19 pneumonia: a double-edged sword. *Radiother Oncol*. 2020.
46. Kirkby C, Mackenzie M. Response to: Low dose radiation therapy for COVID-19 pneumonia a double-edged sword. *Radiother Oncology*. 2020.
47. Jin S, Jiang H, Cai L. New understanding of the low-dose radiation-induced hormesis. *Radiation Medicine and Protection*. 2020.
48. Trott K-R, Kamprad F. Estimation of cancer risks from radiotherapy of benign diseases. *Strahlentherapie und Onkologie*. 2006; 182(8): 431-6.
49. Jansen JTM, Broerse JJ, Zoetelief J, Klein C, Seegenschmiedt HM. Estimation of the carcinogenic risk of radiotherapy of benign diseases from shoulder to heel. *Radiother Oncol*. 2005; 76(3): 270-7.
50. Mazonakis M, Damilakis J. Cancer risk after radiotherapy for benign diseases. *Physica Medica*. 2017; 42: 285-91.
51. Mckeown SR, Hatfield P, Prestwich RJ, Shaffer RE, Taylor RE. Radiotherapy for benign disease; assessing the risk of radiation-induced cancer following exposure to intermediate dose radiation. *Br J Radiol*. 2015; 88(1056): 20150405.
52. Seegenschmiedt M, Micke O, Muecke R, Diseases GCGoRfN-m. Radiotherapy for non-malignant disorders: state of the art and update of the evidence-based practice guidelines. *Br J Radiol*. 2015; 88(1051): 20150080.