

ONLINE FIRSTCASE REPORT**Atrial Septal Defect in an Aging Woman**

Ranjeeta Bai, Sarpat Sardar

ABSTRACT

Atrial Septal Defect (ASD) is a condition that results in an abnormal communication between the right and left atria. In adults, one of the most common types of congenital heart diseases (CHD) is ASD. Around 50 million adults have congenital heart disease (CHD) problems worldwide. This is a report of a 66-year-old female patient diagnosed with ASD around the age of 40 years due to the result of late appearance of signs and symptoms. Usually, the signs and symptoms of ASD develop around 30-40 years. Mostly signs and symptoms may be absent and may be late appearances due to the malformation of the heart. The best control of ASD and related complications and also its death before the age of 25 ASD closure. This case report does not focus only on surgery, angioplasty and catheter-based treatments but also emphasizes that education is very important for these patients. Although life expectancy is not normal but patients generally survive up to adulthood without any surgical intervention, catheter-based treatments but many have reached to advanced age with medication as well as proper rehabilitation.

KEYWORDS: Atrial Septal Defect (ASD), congenital heart disease (CHD), catheter-based treatments, patients' education.

ONLINE FIRST

INTRODUCTION

Atrial Septal Defect (ASD) is a condition that results in an abnormal communication between the right and left atria. ASD is of different types such as; (75%) Secundum ASD, (20%) Ostium primum ASD, (5%) Sinus venosus ASD and (<1%) Coronary ASD. One of the most common types of congenital heart diseases (CHD) in adults is ASD^{1,2}, around 50 million adult have congenital heart disease (CHD) problem worldwide^{3,4}. Several studies shown that CHD has been improved and also decreased mortality rate, Unfortunately, in adult mortality rate still high as compare than general population^{3,4}.

Now day different kinds of treatments are available for ASD adult patients such as surgical, catheter-based treatment in both developed and developing countries. Especially, the Western countries are pressurized and shown that additional efforts to treat properly and control ASD in adolescents as well as adult patients. Unfortunately, the developing countries still have economic issues, poor health services, poor diagnosis and insufficient knowledge related management for ASD patients which leads to increased ASD related issues still¹.

CASE REPORT

A 66 years old female patient diagnosed with Secundum type II with left to right shunt ASD and two Coronary Arteries blockage is reported. When she was born, initially she was asymptomatic but gradually she started facing health problems in her life. With the passage of time, patient's condition gradually becomes poor and she experiences difficulty in breathing during exercise, dry cough and fatigue. At about the age of 40, her family noticed her symptoms and finally she was diagnosed with ASD, surgery was recommended to her cardiologist but unfortunately her husband could not afford the surgery and she was surviving with only medicine.

Around the age of 50 to 55 her children took a decision for her surgery, but she refused believing that surgery might worsen her condition. When she became 60 years old, she got a silent heart attack on 23rd February-2014, and according to cardiologist that silent heart attack was three days old. She was immediately admitted in Isra Hospital, Hyderabad and her investigations were done which included; blood tests, electrocardiogram (ECG), echocardiography (ECHO) and angiography.

ONLINE FIRST

RESULTS

ECG and ECHO was done and result was both interpreted like ECG result was Normal sinus rhythm, Right bundle branch block, Anteroseptal infarct, possibly acute lateral injury pattern and Acute Myocardial infarction and ECHO result was Dilated right sided chambers and left atrium, Moderately reduced left ventricular systolic function with SWMA (segmental wall-motion abnormalities), Mild to moderate mitral regurgitation, Moderate pulmonary artery hypertension and Secundum type ASD with left to right shunt. Fourth day of admission, angiography was done and result was found to be two vessels coronary arterial disease (CAD).

For 5 days she was admitted in Tabba Heart Institute, Karachi and cardiologist prescribed medicines and discharge her from the hospital. After fifteen days she visited the hospital again and her few investigations were done for further treatment. Such as: Electrocardiogram, Echocardiography, transesophageal Echocardiogram and Dobutamine Stress Echocardiography. After investigations, her cardiologist suggested Transcatheter Aortic Valve Implantation (TAVI) but she refused again. Her cardiologist prescribed medicines and some laboratory tests including; HbA1c, potassium, creatinine, GFR and sodium. These tests were repeated every 4 months before her each visit to the hospital.

Prescription

Tablet loprin 75mg 1×OD (once a day), tablet Diamicron MR 30mg 2×OD, tablet Valtec 80mg 1×HS (sleeping hour), tablet lasix 40mg 1×OD, tablet Nebrol fort 1×BD (twice a day), tablet Carvida 6.25mg 1×B, tablet Zopent 40mg 1×OD, tablet Orva 20mg 1×HS and Qalsan-D 1×B.

At present, she goes to hospital for her follow-up every 4 month according to her cardiologist and she is only medicine dependent. Neither any surgery nor any angioplasty is suggested to her by keeping her age factor in view. Her condition is fine but she still has difficulty in breathing during exercise, dry cough and suffocation.

ONLINE FIRST

TABLE I: BLOOD TESTS OF COMPLETE BLOOD COUNT (CBC) OF AGING WOMEN WITH ASD DIAGNOSIS

Complete Blood Count (CBC)	Result	Normal Range
Hemoglobin	11.9gm/dl	11.5-16.0
Haematocrit	36.1%	37-47
Red Blood Cells (R.B.C)	4.05 10e12/L	3.5-5.5
Mean Corpuscular Volume (M.C.V)	89.1 FL	
Mean Corpuscular Hemoglobin (M.C.H)	29.4 PG	27-32
Mean Corpuscular Hemoglobin Concentration (M.C.H.C)	33.0%	30-35
White Blood Cells (W.B.C)	13.0 × 10E9/L	4.0-11.0
Neutrophils	72%	40-75
Lymphocytes	18%	20-45
Eosinophils	02%	1-6
Monocytes	08	2-10
Basophils	00%	<1
Platelet Count	291 × 10E9L	150-400
ESR (Erythrocyte Sedimentation Rate)	36 mm in Hr	0-25
Cholesterol (serum)	179 mg/dl	>200mg

TABLE II: BLOOD TESTS OF RENAL STUDIES OF AGING WOMEN WITH ASD DIAGNOSIS

Renal Studies	Result	Normal Range
Urea (blood)	50 mg/dl	20-45
Creatinine (serum)	0.9 mg/dl	0.7-1.2
Sodium	141 mmol/L	136-149
Potassium	4.5 mmol/L	3.8-5.2
Chloride	102 mmol/L	98-107
Bicarbonate	21 mmol/L	25-29
Uric Acid (serum)	5.6 mg/dl	2.6-6

ONLINE FIRST

TABLE III: BLOOD TESTS OF CARDIAC STUDIES OF AGING WOMEN WITH ASD DIAGNOSIS

Cardiac Blood Tests	Result	Normal Range
(Cardiac Enzymes) KMB	218 U/L	Up to 25
Troponin I	>50.000 ng/ml	Female less than or equal to 0.13 Male less than or equal to 0.33
Brain Natriuretic Peptide (BNP) and N-terminal pro b-type natriuretic peptide (NT-pro BNP)	6725Pg/ml	

DISCUSSION

In adults, around the CHD 22-40% and 10% ASD was recorded. Different types of ASD but most common type of ASD is Secundum ASD⁶. Usually, symptoms appear by age because ASD is asymptomatic in infancy and childhood. About 90% of patients coming in hospital by the age of 40 year experiences of symptoms such as difficulty in breathing during exercise, fatigue, palpitation without treatment^{6,7}. Similarly, this patient also had history of difficulty in breathing during exercise, dry cough and suffocation since 40 years, and now she is 66 years old and patient has multiple complications for example; Ischemic Heart Disease (IHD) with two arteries blocked and myocardial infarction. At present patient is only medicine depended and no any history of angioplasty or surgical treatment is done. Mostly sign and symptoms may be absent and may be late appearances of due to the malformation of the heart^{8,9}. Most children are asymptomatic, though some may experience fatigability very soon and exertional dyspnea. Symptoms usually take about 30-40 years to develop. Nowadays, there are different types of treatments available for closure of ASD such as surgical treatment, catheter-based treatment etc. as well as medicine but health education plays very important role in this kind of patient^{10,11}. Mostly sign and symptoms appear very late age and it's not easy & affordable for all patients such as this patient.

CONCLUSION

The most common congenital disease in adults is ASDS. The best management of ASD can control its related complications and also death before the age of 25. This case report does not focus only on surgery, angioplasty and catheter-based treatments but also emphasizes the education and about ASD. Education plays a very important role in these patients. Although life expectancy is not normal but patients generally survive up to adulthood without any surgical intervention, catheter-based treatments but many have reached to advanced age with medication as well as proper rehabilitation.

Conflict of interest: No any conflict.

Funding: Self-funding.

AUTHOR CONTRIBUTION

Bai R: Created idea, literature searching, and manuscript writing.

Sardar S: Proof reading and critical review

ONLINE FIRST

REFERENCES

1. Xuan Tuan H, Phuoc Long Phan, Duy Kien V, Manh Cuong L, Van Son N, Dalla-Pozza R. (2019). Trends in the prevalence of atrial septal defect and its associated factors among congenital heart disease patients in Vietnam. *Journal of Cardiovascular Development and Disease*, 7(1), 2
2. Rao PS, Harris AD. Recent advances in managing septal defects: atrial septal defects. *F1000Res*. 2017;6:2042.
3. Kurniawaty J, Widyastuti Y. Outcome of adult congenital heart disease patients undergoing cardiac surgery: clinical experience of dr. Sardjito hospital. *BMC Proc*. 2019;13(Suppl 11):16.
4. Engelings CC, Helm PC, Abdul-Khaliq H, Asfour B, Bauer UMM, Baumgartner H, et al. Cause of death in adults with congenital heart disease - An analysis of the German National Register for Congenital Heart Defects. *Int J Cardiol*. 2016;211:31–6.
5. Płońska-Gościniak E, Department of Cardiology Pomeranian Medical University, Szczecin, Poland, Kukulski T, Kasprzak JD, Gąsior Z, Szyszka A, et al. Stress echocardiography. Part II: Stress echocardiography in conditions other than coronary heart disease. *J Ultrason*. 2019;19(76):49–53.
6. Sen SS, Barua T, Dey D, Chowdhury MA, Nessa L. Pattern of congenital Heart Disease in children presenting at paediatric cardiology unit in chattagram maa shishu-O-general hospital, Chittagong. *Chattagram Maa-O-Shishu Hosp Med Coll j*. 2018;16(2):40–3.
7. Yang M-C, Wu J-R. Recent review of transcatheter closure of atrial septal defect. *Kaohsiung J Med Sci*. 2018;34(7):363–9.
8. Bissessor N. Current perspectives in percutaneous atrial septal defect closure devices. *Med Devices (Auckl)*. 2015;8:297–303.
9. Wang SY, Welch TD, Eifenbein A, Kaplan AV. Spontaneous closure of a secundum atrial septal defect. *Methodist Debaque Cardiovasc J*. 2018;14(1):60–2.
10. Oster M, Bhatt AB, Zaragoza-Macias E, Dendukuri N, Marelli A. Interventional therapy versus medical therapy for secundum atrial septal defect: A systematic review (part 2) for the 2018 AHA/ACC guideline for the management of adults with congenital heart disease: A report of the American college of cardiology/American heart association task force on clinical practice guidelines: A report of the American college of cardiology/American heart association task force on clinical practice guidelines. *Circulation*. 2019;139(14):e814–30.
11. Diaconu CC. Atrial septal defect in an elderly woman-a case report. *J Med Life*. 2011;4(1):91–3.