

ORIGINAL ARTICLE

**Frequency of Asymptomatic Bacteriuria in Patients
with Preterm Labor**

Samana Aleem, Hazooran Lakhan, Shahida Karamat, Asifa Abdul Jabbar, Saima Farook,
Farheen Shaikh

Dr. Samana Aleem

Senior Registrar
PUMHS Nawabshah, Sindh-Pakistan.

Dr. Hazooran Lakhan

Senior Registrar
Ghulam Muhammad Mahar Medical College
Sukkur, Sindh-Pakistan.

Dr. Shahida Karamat

Assistant Professor
Bahria University Medical and Dental College
Islamabad-Pakistan.

Dr. Asifa Abdul Jabbar (*Corresponding Author*)

Assistant Professor
Department of Obstetrics & Gynaecology
Gambat Institute of Medical Sciences (GIMS)
Khairpur Mirs, Safeer Autos, Agra Road, Gambat, Sindh-Pakistan.
Email: drasifaabduljabbar@gmail.com

Dr. Saima Farooq

Consultant
Department of Obstetrics & Gynecology
Hamdard Hospital, Karachi, Sindh-Pakistan.

Dr. Farheen Sheikh

Assistant Professor
Baqai Medical University, Karachi, Sindh-Pakistan.

ABSTRACT

OBJECTIVE: To determine the frequency of asymptomatic bacteriuria in patients presenting in preterm labor in a tertiary care hospital

METHODOLOGY: This Cross-sectional study was conducted at Department of Obstetrics & Gynecology, Peer Syed Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat from July to December 2018. A total of 115 women with preterm labour were included in this study through Non probability consecutive sampling technique. Clean voided mid-stream urine specimen of patients was collected. Subjects were adequately educated to prevent contamination of specimen. The specimen was collected into sterilized wide, necked, leak proof, plastic containers. The patients were followed for outcome of preterm labour. SPSS v. 20 was used to enter and analyses the data.

RESULTS: The average age of patients was 31.94 ± 5.26 years. The mean gestational age at delivery was 32 ± 3.61 weeks. Frequency of asymptomatic bacteriuria was 27.83%.

CONCLUSION: It has been concluded that a rate of asymptomatic bacteriuria can be observed among pregnant females admitted with preterm labour. The hazardous consequences related to asymptomatic bacteriuria can be minimized by prescribing a course of antimicrobials in early pregnancy.

KEY WORDS: Urinary tract infection, asymptomatic bacteriuria, Preterm labor

INTRODUCTION

Urinary tract infection (UTI), a serious health problems affecting millions of people each year accounting for about 8.3 million visits to the hospital each year^{1,2}. The spectrum ranges from asymptomatic bacteria to severe pyelonephritis¹. It is the second most common type of infections in the body. The common cause of detected usually in uncomplicated UTI are *E. coli* (75 – 90%), *staphylococcus saprophyticus* (5 – 15 %), *Enterococcus* and other gram negative rods other than *E. coli* have also been implicated in some cases². Studies have also found increased infections with *E.coli* and *Staphylococcus* in patients with UTI³.

Asymptomatic bacteriuria may proceed to symptomatic UTI and is responsible for over 6 million outpatient visits each year with a prevalence of 4-7% in pregnancy⁴. As it is evident from previous studies that asymptomatic bacteriuria developed in the pregnancy considerably increases the chances of low birth weight as well as significant number of preterm deliveries⁵. The prevalence of asymptomatic bacteriuria is higher (25.68%) in third trimester than first trimester (15.79%)⁶.

Thus early identification of asymptomatic bacteriuria is of utmost importance as it has potential for saving pregnant women, in whom bacteriuria is an established risk factor for serious complication including the endometritis, chorioamnionitis, reduced fetal growth, preterm labour & birth, raised still-birth or perinatal mortality, mental retardation and slow growth^{2,7}. A large population-based study in Israeli women revealed 2.5% rate of asymptomatic bacteriuria and 2.3% rate of symptomatic bacteriuria. Mazor et al., states that findings are indicators for the strength of the prenatal care instead of the precise causal effects of the UTI⁸. Similarly, few surveys related to the different races established that the variations among the females belong to the Native America and observed the highest prevalence of such urinary infections was 24.2% among Native American females than females in Asia (10.3%), England (16.6%), Spain (18.3%), and African countries (20.3%)^{9,10}. Therefore, the aim of our study is to determine the frequency of asymptomatic bacteriuria in patients with preterm labor if results of study show high prevalence of asymptomatic bacteriuria in preterm labour so that efforts should be made for its earlier control to reduce prenatal morbidity and mortality.

METHODOLOGY

This cross-sectional study was done at Department of Obstetrics & Gynecology at Peer Syed Abdul Qadir Shah Jeelani Institute of Medical Sciences, Gambat from July to December 2018. By taking the frequency of asymptomatic bacteriuria among women having preterm labour is 25.68% confidence level 95% from the margin of error 8% and at least sample of 115 were included through Non probability consecutive sampling technique. Pregnant females aged 20 – 45 years presented in preterm labor (labor and delivery between 24-36 weeks of pregnancy) were included. Females who were already diagnosed with UTI and taking antibiotics, multiple gestation, polyhydramnios, had cervical surgery or uterine anomaly were excluded from the study. Written informed consent was obtained from pregnant women who fulfill inclusion criteria. Clean voided mid-stream urine specimen of patients was collected. Subjects were adequately educated to prevent contamination of specimen. The specimen was collected into sterilized wide, necked, leak proof, plastic containers. The patients were followed for outcome of preterm labour. Asymptomatic bacteriuria was labelled if there was a significant bacterial count $> 10^5$ CFU/ml present in the urine of a person without symptoms. After collection of data, the analysis was done using SPSS software version 20. Frequency was calculated for asymptomatic bacteriuria. Stratification with respect to age and parity was done. Post stratification, Chi-square test was applied with p-value ≤ 0.05 as significant.

RESULTS

The average age of the patients was 31.94±5.26 years similarly mean gestational age was 32±3.61 weeks. Out of 115 women, 41(35.56%) were primiparous and 71(64.35%) were multiparous. Table I.

Frequency of the asymptomatic bacteriuria among patients presenting in preterm labor was 32 (27.83%). Figure I.

Rate of asymptomatic bacteriuria among patients presenting in preterm labor was insignificant among different age groups ($p = 0.32$) as well as in parity strata ($p > 0.05$). Table II.

TABLE I: BASELINE CHARACTERISTICS OF PATIENTS

n	115
Age (Years)	31.94±5.26
Gestational Age (Weeks)	32±3.61
Primiparous	41 (35.7%)
Multiparous	74 (64.4%)

FIGURE I: FREQUENCY OF ASYMPTOMATIC BACTERIURIA IN PATIENTS WITH PRETERM LABOR




TABLE II: COMPARISON OF ASYMPTOMATIC BACTERIURIA IN DIFFERENT AGE AND PARITY GROUPS

	Asymptomatic Bacteriuria		p-value
	Yes	No	
Age (years)			0.32
21-25	2 (6.25%)	13 (15.66%)	
26-30	9 (28.13%)	18 (21.69%)	
31-35	11 (34.38%)	35 (42.17%)	
> 35	10 (31.25%)	17 (20.48%)	
Parity			0.85
Primiparous	11 (34.38%)	30 (36.14%)	
Multiparous	21 (65.63%)	53 (63.86%)	

DISCUSSION

Asymptomatic bacteriuria can be present in 2-10% pregnancies.¹¹ In several observational studies, conducted in developing countries, the rate of symptomatic bacteriuria among pregnant women can be from 4% to 10%^{5,12-14}. This disparity among different studies may be due to numerous factors e.g., geographical deviation, socioeconomic status, ethnicity, level of healthcare center (primary care, community based or hospital), and screening tests distinction (urinary dipstick, microscopy or culture). There is a significant disparity in rates of bacteriuria in different races, besides there is also difference in rates in same race due to different geographical living regions or socioeconomic status. Symptomatic bacteriuria does not cause much problems as it is easy to diagnose and treat because of its obvious symptoms. However asymptomatic bacteriuria is problematic to diagnose and it is more prevalent during pregnancy¹⁵.

It is specifically important during pregnancy as 30-40% untreated cases of asymptomatic bacteriuria can develop acute pyelonephritis in later trimesters of pregnancy¹⁶⁻¹⁸. Fetomaternal complications of asymptomatic bacteriuria are suggestive urinary tract infections, pyelonephritis, toxemia related to preeclampsia, anemia, low birthweight, intrauterine growth retardation, premature rupture of membrane, preterm labour or endometritis¹⁹⁻²¹.

Though, in developed countries, the evaluation and management of asymptomatic bacteriuria during first trimester of pregnancy is standard-of-care and role of precise antibacterial treatment during pregnancy is well known²². Our study reported that the out of 115 women with preterm labour, frequency of asymptomatic bacteriuria was 27.83%. The stated frequency of asymptomatic bacteriuria in pregnancy was were 30% in Yemen,²³ 4.8% in United Arab Emirates²⁴, 9.9% in Qatar²⁵ & 3.3 - 6.1% in Iran²⁶. Now-a-days, the association of asymptomatic bacteriuria with obstetrical-related complication particularly preterm delivery is a recognized fact^{27,28}.

But the physiology of association between asymptomatic bacteriuria and preterm labour is not well-established yet, but a hypothetical dispute is made for contributory role for secretion of phospholipase A2 by pathogens, which can then cause an early initiation of labor by stimulation of prostaglandin²⁹. Many researchers have detected the high rate of pyelonephritis among pregnant females having bacteriuria³⁰.

It is conclusive that the impact of bacteriuria on preterm labor might be indirectly facilitated by pregnancy induced hypertension. It is also probable that bacteriuria can directly affect preterm labor, by amnionitis development. It was heretofore proposed that bacterial infection in amniotic fluid can be a risk factor of preterm delivery^{24,31}.

CONCLUSION

It has been concluded that there is a high prevalence of asymptomatic bacteriuria among pregnant females presented in labour during preterm weeks of gestation. The adverse outcomes caused by asymptomatic bacteriuria during pregnancy can be decreased by using antibiotic therapy earlier in the pregnancy. Thus, evaluation and cure of asymptomatic bacteriuria is required to be integrated in routine antenatal care for cohesive approach to safe the mother and neonate without compromising their health and life.

Ethical permission: Pir Abdul Qadir Shah Jeelani Institute of Medical Sciences Gambat, REC letter No. D.PAQSJIMS.REC/UTI/2018/200, Dated 20-7-2018.

Conflict of Interest: There is no conflict of interest.

Funding: There was no any funding agency.

AUTHOR CONTRIBUTIONS

Aleem S: Data collection

Lakhan H: Data collection

Karamat S: Data analysis

Jabbar AA: Result formulation

Farook S: Result formulation

Shaikh F: Paper formulation

REFERENCES

1. Assafi MS, Ibrahim NM, Hussein NR, Taha AA, Balatay AA. Urinary bacterial profile and antibiotic susceptibility pattern among patients with urinary tract infection in duhok city, kurdistan region, Iraq. *Int J Pure Appl Sci Technol*. 2015; 30(2): 54.
2. Kline KA, Lewis AL. Gram-positive uropathogens, polymicrobial urinary tract infection, and the emerging microbiota of the urinary tract. *Microbiol Spectr*. 2016; 4(2).
3. Saha S, Nayak S, Bhattacharyya I, Saha S, Mandal A, Chakraborty S, et al. Understanding the patterns of antibiotic susceptibility of bacteria causing urinary tract infection in West Bengal, India. *Front Microbiol*. 2014; 5: 463.
4. Cohen R, Gutvitz G, Wainstock T, Sheiner E. Maternal urinary tract infection during pregnancy and long-term infectious morbidity of the offspring. *Early Hum Dev*. 2019; 136: 54-9.
5. Matuszkiewicz-Rowińska J, Małyszko J, Wieliczko M. Urinary tract infections in pregnancy: old and new unresolved diagnostic and therapeutic problems. *Arch Med Sci*. 2015; 11(1): 67.
6. Oli AN, Okafor CI, Ibezim EC, Akujiobi CN, Onwunzo MC. The prevalence and bacteriology of asymptomatic bacteriuria among antenatal patients in Nnamdi Azikiwe University Teaching Hospital Nnewi; South Eastern Nigeria. *Nigerian J Clin Pract*. 2010; 13(4): 409-12.
7. Chen Y, Huang L, Zhang H, Klebanoff M, Yang Z, Zhang J. Racial disparity in placental pathology in the collaborative perinatal project. *Int J Clin Exp Pathol*. 2015; 8(11): 15042.
8. Mazor-Dray E, Levy A, Schlaeffer F, Sheiner E. Maternal urinary tract infection: is it independently associated with adverse pregnancy outcome? *J Mat Fet Neonat Med*. 2009; 22(2): 124-8.
9. Dieke AC, Zhang Y, Kissin DM, Barfield WD, Boulet SL. Disparities in Assisted Reproductive Technology Utilization by Race and Ethnicity, United States, 2014: A Commentary. *Journal of women's health (2002)*. 2017; 26(6): 605-8.
10. Harris DE, Aboueissa A-M, Baugh N, Sarton C, Lichter E. 11-year trends in pregnancy-related health indicators in maine, 2000–2010. *J Pegnan*. 2014; 2014.
11. Smaill FM, Vazquez JC. Antibiotics for asymptomatic bacteriuria in pregnancy. *Cochrane Database Syst Rev*. 2015; (8): CD000490
12. Sujatha R, Nawani M. Prevalence of asymptomatic bacteriuria and its antibacterial susceptibility pattern among pregnant women attending the antenatal clinic at Kanpur, India. *J Clin Diag Res*. 2014; 8(4): DC01.
13. Bernabe KJ, Langendorf C, Ford N, Ronat J-B, Murphy RA. Antimicrobial resistance in West Africa: a systematic review and meta-analysis. *Int J Antimicrob Agent*. 2017; 50(5): 629-39.
14. Glaser AP, Schaeffer AJ. Urinary tract infection and bacteriuria in pregnancy. *Urol Clin*. 2015; 42(4): 547-60.
15. Shaheen H, Farahat T, El-Hakeem Hammad N. Prevalence of urinary tract infection among pregnant women and possible risk factors. *Menoufia Med J*. 2016; 29(4): 1055-9.
16. El-Sokkary M. Prevalence of Asymptomatic Bacteriuria in Antenatal Women with Preterm Labor at an Egyptian Tertiary Center. *J Am Sci*. 2011; 7(4): 605-10.
17. Kalinderi K, Delkos D, Kalinderis M, Athanasiadis A, Kalogiannidis I. Urinary tract infection during pregnancy: current concepts on a common multifaceted problem. *J Obstet Gynaecol*. 2018; 38(4): 448-53.

18. Mukherjee M, Koley S, Mukherjee SK, Basu S, Ghosh B, Chakraborty S. Phylogenetic background of *E. coli* isolated from asymptomatic pregnant women from Kolkata, India. *J Infect Develop Countries*. 2015; 9(07): 720-4.
19. Faidah HS, Ashshi AM, El-Ella GAA, Al-Ghamdi AK, Mohamed AM. Urinary tract infections among pregnant women in Makkah, Saudi Arabia. *Biomed Pharmacol J*. 2015; 6(1): 01-7.
20. Ortega-González Y, Castro-Diaz D. Antibiotic considerations for urinary tract infections in pregnancy. *Curr Bladder Dysfunct Rep*. 2014; 9(3): 167-74.
21. Szweda H, Józwiak M. Urinary tract infections during pregnancy-an updated overview. *Dev Period Med*. 2016; 20(4): 263-72.
22. Vazquez JC, Abalos E. Treatments for symptomatic urinary tract infections during pregnancy. *Cochrane Database Syst Rev*. 2011; (1): Cd002256.
23. Al-Haddad AM. Urinary tract infection among pregnant women in Al-Mukalla district, Yemen. *East Mediter Health J*. 2005; 11(3): 505-10.
24. Abdullah AA, Al-Moslih MI. Prevalence of asymptomatic bacteriuria in pregnant women in Sharjah, United Arab Emirates. *East Mediter Health J*. 2005; 11(5-6): 1045-52.
25. Al-Kuwari M, Taher Aseel M, Mohamed Al-Meer F, F Ismail M. Prevalence and Predictors of Asymptomatic Bacteriuria among Pregnant Women Attending Primary Health Care in Qatar. *Middle East J Fam Med*. 2009; 7: 10.
26. Jazayeri Moghadas A, Irajian G. Asymptomatic Urinary Tract Infection in Pregnant Women. *Iran J Pathol*. 2009; 4(3): 105-8.
27. Misra R. *Ian Donald's Practical Obstetric Problem*. 6th ed. India: B.I. Publications Pvt. Limited; 2006.
28. Gharbi M, Drysdale JH, Lishman H, Goudie R, Molokhia M, Johnson AP, et al. Antibiotic management of urinary tract infection in elderly patients in primary care and its association with bloodstream infections and all cause mortality: population based cohort study. *Br Med J*. 2019; 364: 1525.
29. Romero R, Miranda J, Chaiworapongsa T, Korzeniewski SJ, Chaemsaitong P, Gotsch F, et al. Prevalence and Clinical Significance of Sterile Intra-amniotic Inflammation in Patients with Preterm Labor and Intact Membranes. *Am J Reprod Immunol*. 2014; 72(5): 458-74.
30. Imade PE, Izekor PE, Eghafona NO, Enabulele OI, Ophori E. Asymptomatic bacteriuria among pregnant women. *N Am J Med Sci*. 2010; 2(6): 263-6.
31. Tugrul S, Oral O, Kumru P, Köse D, Alkan A, Yildirim G. Evaluation and importance of asymptomatic bacteriuria in pregnancy. *Clin Exp Obstet Gynecol*. 2005; 32(4): 237-40.