

ORIGINAL ARTICLE

MRI Grading of Lumbar Spine Degenerative Disc Disease using a Modified Pfirrmann Grading System

Rabia Hafeez, Inayatullah Memon

Dr. Rabia Hafeez

Senior Registrar
Department of Radiology
Ghulam Muhammad Mahar Medical College
Sukkur, Sindh-Pakistan.
Email: rabiahafeez66@gmail.com

Dr. Inayatullah Memon

Assistant Professor
Department of Radiology
Ghulam Muhammad Mahar Medical College
Sukkur, Sindh-Pakistan.

Ahead of Print

ABSTRACT

OBJECTIVE: To evaluate the extent of involvement and severity of intervertebral degenerative disc disease on MRI lumbar spine by using a modified pfirrmann grading system in patients with low back pain.

METHODOLOGY: This cross-sectional study was conducted at the radiology department of GMC hospital Sukkur from August 2021- January 2022. T2 weighted images of the MRI lumbar spines of 460 consecutive patients were retrospectively reviewed and the severity of degeneration was assessed using a modified pfirrmann grading system. Patients with imaging features of degenerative disc disease on MRI lumbar spine were included in the study. Patients with a history of trauma, surgery and infection of the lumbar spine were excluded from the study. Data was collected and analyzed using SPSS version 21.

RESULTS: Disc degeneration was classified by a modified pfirrmann grading system score from 1-8, with no disc score in grades 1, 2 and 3. Other discs score were 13(2.82%), 41(8.9%), 187(40.6%), 125(27.1%) and 94(20.4%) in grade 4,5,6,7 and 8 respectively.

CONCLUSION: This study concludes that a modified pfirrmann grading system is helpful in the assessment of the severity of lumbar spine intervertebral disc degeneration. It is an easy and reliable method and can be easily applicable for better correlation between radiological findings and clinical diagnosis. This grading helps in early detection and decides the mode of treatment. At present lack of precise knowledge and data available in this context in our country.

KEYWORDS: MRI lumbar spine, low back pain, intervertebral disc degeneration, intervertebral disc scoring system, modified pfirrmann grading.

INTRODUCTION

Lumbar intervertebral disc degeneration is considered a significant risk factor for lower back pain besides reducing bone density, muscular spasm and spondylolisthesis and becoming a significant health issue^{1,2}.

Disc herniation, spinal canal stenosis and facet joint arthrosis are different forms of degenerative disc disease of the lumbar spine³. Protrusion of the central part of the disc due to disc thinning and annular tear results in disc herniation. Displacement of the annulus fibrosus outer border outside the outlines of the vertebral body represents disc bulge⁴. Narrowing the spinal column, lateral recess, or neural foramina refers to spinal canal stenosis that causes nerve root compression and low back pain^{5,6}.

On the T2 W sequence of MRI, the degenerative changes in the disc appear as a gradual loss of standard high signal of the nucleus pulposus⁷. Pfirrmann CW 2001⁷ proposed a grading system and classification based on signal intensity of disc morphology, differentiation between annulus fibrosus and nucleus pulposus and height of the disc, labelled as grade I-V^{8,9}. Another system of grading was modified Pfirrmann system assigns grades 1-8¹⁰.

The modified Pfirrmann grading system is a convenient and valuable semi-quantitative evaluation of disc degeneration with good reproducibility¹¹.

In this study, we evaluate the extent of involvement and severity of intervertebral degenerative disc disease on MRI lumbar spine using a modified pfirrmann grading system.

METHODOLOGY

This cross-sectional study was performed at the Radiology department of Ghulam Muhammad Mahar medical college hospital Sukkur from August 2021-January 2022. Non-probability consecutive sampling technique was used. After taking informed consent Total of 460 cases with imaging features of degenerative disc disease on MRI lumbar spine were included in the study. Patients with a history of trauma, surgery and infection of the lumbar spine were excluded from the study.

MRI lumbar spine screening includes T2 axial and sagittal sequences obtained on a 0.35 Tesla open MRI machine. Two independent radiologists reviewed T2 sagittal sequences with five-year experience.

The intervertebral disc signal intensity changes were assessed at each level on the sagittal T2 weighted sequence of MRI lumbar spine using Modified Pfirrmann grading systems. Results were expressed as grades 1-8. Data was collected and analyzed using SPSS version 21.

Ahead of Print

RESULTS

Of 460 MRI lumbar spine T2 sagittal images, 267(58%) were male, and 193(42%) were female. The ages of patient range from 25-65 years. The mean age was 45 years, and the most common age group was 30-55.

Degenerated lumbar spine discs were graded using a modified pfirrmann Grading system.

Grading by modified pfirrmann shows no disc score in grades 1, 2 and 3. Other discs score 13(2.82%), 41(8.9%), 187(40.6%), 125(27.1%) and 94(20.4%) in grade 4,5,6,7 and 8 respectively. **Table I**

TABLE I: MODIFIED PFIRRMANN GRADES OF THE LUMBAR SPINE ON MRI

MODIFIED PFIRRMANN GRADES	Score (%)
I	0
II	0
III	0
IV	13(2.82%)
V	41(8.91%)
VI	187(40.65%)
VII	125(27.17%)
VIII	94(20.43%)
Total	460(100%)

DISCUSSION

At different ages, low back pain is the primary health issue affecting two-thirds or 50-80% of the population worldwide^{12,13}. MRI is considered a baseline radiological investigation with better resolution and lack of radiation in diagnosing lumbar spine degeneration, and it allows early identification and thus prevents long-term neurological complications^{14,15}.

Another benefit of MRI is better identifying soft tissue detail like discs, nerves and para-spinal muscles responsible for causing low back pain¹⁶.

Pfirschmann grading systems score the signal intensity and morphology of the disc on T2 weighted sagittal sequence into five grades. Grades I and II saw as average central bright signal intensity. Grade III appears as a disc with decreased signal intensity. Grade IV and V show increasing disc height loss and additional degenerative changes. Grade III disc labelled as dehydrated or desiccated is of clinical importance as it depicts early changes and is suitable for less invasive medical treatment. This grading distinguishes it from grade IV, which needs surgical intervention^{17,18}.

Kettler A 2006¹⁹ reviewed that the modified pfirschmann grading system was the only MR grading system that showed good inter and intra reader reliability with kappa values of > 0.6.

Grading and incidence of disc degeneration were affected by age and sex in older age groups and is a natural phenomenon after 30 years²⁰. Pfirschmann grade IV (Moderate disc degeneration) in lower lumbar levels predisposes herniation of disc and indication of surgery after spinal stenosis²¹. Schneiderman G et al.²³ proposed a modified grading system for better evaluation of degeneration of intervertebral disc comprised of 1-8 grades.

Grades 1-3 build on the intensity of signals returning from the nucleus pulposus and inner fibres of the annulus. Grades 4 and 5 showed indistinct borders between the outer and inner fibres of the posterior annulus. Grade 6-8 shows, in addition, progressive reduced disc height¹⁰.

MRI is considered the best imaging modality for diagnosing and evaluating disc degeneration in the lumbar spine. Quantitative classification can also be performed according to the changes in signal in the nucleus pulposus²². The first classification of disc degeneration was Schneiderman G et al.²³ based on indications of intervertebral discs on T2-weighted sequence classified into four grades. Quantitative classification of relative signal intensity (RSI) of nucleus pulposus was utilized by Luoma K 2001²⁴ and his colleagues in their study.

Our study results were similar to studies done by Yu LP 2012²⁵ shows Modified pfirschmann grades scores were 1.85%, 14.82%, 39.81%, 29.63%, and 13.89% for grades 4, 5, 6, 7 and 8, no grade score for 1-3. Another study shows similar results by Alkhasawneh MH 2021²⁶.

CONCLUSION

This study concludes that the Modified pfirrmann grading system is helpful in the assessment of the severity of lumbar spine intervertebral disc degeneration. It is an easy and reliable method and can be easily applicable for better correlation between radiological findings and clinical diagnosis. This grading helps in early detection and decides the mode of treatment. Lack of precise knowledge and data available in this context in our country.

Ethical Permission: Ghulam Muhammad Mahar Medical College Sukkur, ERC letter No. 121/22/ERC/04, dated 10.08-2021.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure / Grant Approval: No funding agency was used for this research.

Data Sharing Statement: The data supporting this study's findings are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

AUTHOR CONTRIBUTIONS

Hafeez R: Manuscript Writing, Data Collection, Analysis

Memon I: Data collection

REFERENCES

1. Hansen BB, Bendix T, Grindsted J, Bliddal H, Christensen R, Hansen P et al. Effect of Lumbar Disc Degeneration and Low-Back Pain on the Lumbar Lordosis in Supine and Standing: A Cross-Sectional MRI Study. *Spine (Phila Pa 1976)*. 2015; 40(21): 1690-6.
2. Folkvardsen S, Magnussen E, Karppinen J, Auvinen J, Larsen RH, Wong C et al. Does elite swimming accelerate lumbar intervertebral disc degeneration and increase low back pain? A cross-sectional comparison. *Eur Spine J*. 2016; 25(9): 2849-55.
3. Saleem S, Aslam HM, Rehamn MA, Raees A, Alvi AA, Ashraf J. Lumbar disc degenerative disease: disc degeneration symptoms and MRI findings. *Asian Spine J*. 2013; 7(4): 322-24.
4. Wong AY, Karppinen J, Samartzis D. Low back pain in older adults: risk factors, management options, and future directions. *Scoliosis Spinal Disorders*. 2017; 12 (1): 1-23.
5. Arya RK. Low back pain - signs, symptoms, and management. *J Ind Acad Clin Med*. 2014; 15 (1): 30-41.
6. Casser HR, Seddigh S, Rauschmann M. Acute Lumbar Back Pain. *Dtsch Arztebl Int*. 2016; 113(13): 223-34.
7. Pfirrmann CW, Metzdorf A, Zanetti M, Hodler J, Boos N. Magnetic resonance classification of lumbar intervertebral disc degeneration. *Spine (Phila Pa 1976)*. 2001; 26: 1873-78.
8. Beack JY, Chun HJ, Bak KH, Choi KS, Bae IS, Kim KD. Risk Factors of Secondary Lumbar Discectomy of a Herniated Lumbar Disc after Lumbar Discectomy. *J Korean Neurosurg Soc*. 2019; 62(5): 586-93.
9. Farshad-Amacker NA, Farshad M, Winklehner A, Andreisek G. MR imaging of degenerative disc disease. *Eur J Radiol*. 2015; 84(9): 1768-76.
10. Griffith JF, Wang YX, Antonio GE, Choi KC, Yu A. Modified Pfirrmann grading system for lumbar intervertebral disc degeneration. *Spine*. 2007; 32: 708–12.
11. Miyazaki M, Hong SW, Yoon SH, Morishita Y, Wang JC. Reliability of a magnetic resonance imaging-based grading system for cervical intervertebral disc degeneration. *J Spinal Disord Tech*. 2008; 21(4): 288-92.
12. Ramdas J, Jella V. Prevalence and risk factors of low back pain. *Int J Adv Med*. 2018; 5(5): 1120-23.
13. Fatoye F, Gebrye T, Odeyemi I. Real-world incidence and prevalence of low back pain using routinely collected data. *Rheumatic Int*. 2019; 39(4): 619- 26.
14. Cicero G, D'Angelo T, Racchiusa S. Cross-sectional imaging of parotid gland nodules: A brief practical guide. *J Clin Imaging Sci*. 2018; 8(1): 14.
15. Yousof M, Din RU, Ahmad M, Khattak S, Ali S. The Prevalence of Pathologies Causing Low Back Pain Using Magnetic Resonance Imaging for Categorization. *OMICS J Radiol*. 2021; 10: 323. doi: 10.4172/2167-7964.1000323.
16. Tonosu J, Oka H, Higashikawa A, Okazaki H, Tanaka S, Matsudaira K. The associations between magnetic resonance imaging findings and low back pain: A 10-year longitudinal analysis. *PLoS One*. 2017; 12(11): e0188057.
17. Lotz JC, Haughton V, Boden SD, An HS, Kang JD, Masuda K, et al. New treatments and imaging strategies in degenerative disease of the intervertebral disks. *Radiology*. 2012; 264(1): 6-19.
18. Haughton V. Don't you call me desiccated, J. Wellington Wimpy. *AJNR Am J Neuroradiol*. 2010; 31(9): 1551.
19. Kettler A, Wilke HJ. Review of existing grading system for cervical or lumbar disc and facet joint degeneration. *Eur Spine J*. 2006; 15(6): 705-18.
20. Oh CH, Yoon SH. Whole Spine Disc Degeneration Survey according to the Ages and Sex Using Pfirrmann Disc Degeneration Grades. *Korean J Spine*. 2017; 14(4): 148-54.
21. Hwang HJ, Park HK, Lee GS, Heo JY, Chang JC. Predictors of Reoperation after Microdecompression in Lumbar Spinal Stenosis. *Korean J Spine*. 2016; 13(4): 183-89.

22. Xiao L, Ni C, Shi J, Wang Z, Wang S, Zhang J et al. Analysis of Correlation Between Vertebral Endplate Change and Lumbar Disc Degeneration. *Med Sci Monit.* 2017; 15(23): 4932-38.
23. Schneiderman G, Flannigan B, Kingston S, Thomas J, Dillin WH, Watkins RG. Magnetic resonance imaging in the diagnosis of disc degeneration: correlation with discography. *Spine (Phila Pa 1976).* 1987; 12(3): 276-81.
24. Luoma K, Vehmas T, Riihimaki H, Raininko R. Disc height and signal intensity of the nucleus pulposus on magnetic resonance imaging as indicators of lumbar disc degeneration. *Spine (Phila Pa 1976).* 2001; 26(6): 680–86.
25. Yu LP, Qian WW, Yin GY, Ren YX, Hu ZY. MRI assessment of lumbar intervertebral disc degeneration with lumbar degenerative disease using the Pfirrmann grading systems. *PLoS One.* 2012; 7(12): e48074.
26. Alkhasawneh MH, Al-Mnayyis A, Bagain Y. Spinal Degeneration and Degenerative Disc Disease correlation identified with Magnetic Resonance Imaging. *Biomed Pharmacol J.* 2021; 14(1): 491-96.

Ahead of Print