

Effectiveness of the Kampung Bunga Program in Accelerating Stunting Reduction in Central Lampung Regency, Indonesia

Sri Indra Trigunarto^{1*}, Ika Oktaviani², Arie Nugroho³, Yustin Nurkhoiriyah⁴, Febrina Sarlinda⁴

ABSTRACT

OBJECTIVE: To evaluate the association of the Kampung Bunga program with accelerating stunting reduction in Central Lampung Regency.

METHODOLOGY: A cross-sectional study was conducted from May 2025 to February 2026 involving 563 children aged 0–59 months in Central Lampung Regency. The stunting group comprised 138 children, while the non-stunting group comprised 425 children. The sample was recruited using a multistage cluster sampling technique. We used anthropometric measurements, structured questionnaires, and maternal health records for data collection. We measured and calculated clinical nutritional indicators, including height-for-age z-scores (HAZ), weight-for-age z-scores (WAZ), and mid-upper arm circumference (MUAC). Maternal-related risk factors included education level, antenatal care (ANC) attendance, exclusive breastfeeding practices, and nutritional status during pregnancy. Community health intervention variables included participation in Kampung Bunga program activities. Chi-square tests, independent t-tests, and multivariate logistic regression were performed as appropriate for the data, with statistical significance set at $P < 0.05$.

RESULTS: The prevalence of stunting was lower among program participants (18.3%) than among the non-participants (31.7%; $P < 0.001$). Multivariate analysis found that Kampung Bunga participation (AOR = 0.42; 95% CI: 0.28–0.63), complete ANC visits (AOR = 0.51; 95% CI: 0.34–0.76), exclusive breastfeeding (AOR = 0.47; 95% CI: 0.31–0.71), and higher maternal education were identified as significant protective factors against stunting ($P < 0.05$).

CONCLUSION: Stunting prevalence was significantly lower among participants of the integrated community-based nutritional intervention. We recommend scaling up this intervention alongside continued strengthening of maternal education and antenatal care services to accelerate stunting reduction.

KEYWORDS: Stunting, Community Health Intervention, Maternal Risk Factors, Nutritional Indicators

INTRODUCTION

Stunting, defined as a height-for-age z-score standard deviation from the World Health Organization (WHO) Child Growth Standards median, is one of the most prevalent forms of chronic malnutrition worldwide^{1,2}. This status indicates that the child has suffered too long from poor nutrition, repeated infections and inadequate psychosocial stimulation in the first 1,000 days of life³. Worldwide, there are estimated to be 149 million stunted children under five years of age, the vast majority living in Asia and Africa⁴. Indonesia has one of the highest prevalences of childhood stunting in Southeast Asia, with the most

recent estimates indicating that two out of five children suffer from chronic malnutrition⁵. National survey data suggest that the prevalence of stunting in Indonesia remains a serious public health problem at 29.6%, well above the World Health Organization (WHO) limit⁶.

Important regional disparities exist within Indonesia, with rural areas consistently showing a higher prevalence of stunting than urban settings. Lampung Province, located in South Sumatra, has experienced ups and downs in its stunting rate, and several districts have seen their number of stunted children rise sharply over the past few years. Central Lampung Regency, whose population is dominated by rural communities, is classified as a stunting priority area for intervention.

Stunting has a multifactorial etiology with immediate, underlying, and structural determinants⁷. At the immediate level, low dietary intake and infectious diseases directly impede linear growth and development among children⁸. Root causes include household food insecurity, inappropriate feeding and care practices for mothers and young children, and limited access to health services and sanitation⁹. Maternal-related factors have always been shown to be one of the most important predictors of child

¹Poltekkes Kemenkes Tanjung Karang, Tanjung Karang, Lampung, Indonesia

²Department of Midwifery, Poltekkes Kemenkes Tanjung Karang, Tanjung Karang, Lampung, Indonesia

³Department of Nutrition, Poltekkes Kemenkes Tanjung Karang, Tanjung Karang, Lampung, Indonesia

⁴Medical Laboratory of the Department of Technology, Poltekkes Kemenkes Tanjung Karang, Tanjung Karang, Lampung, Indonesia

Correspondence: sriindratrigunarto1@gmail.com
doi: 10.22442/jlumhs.2026.01497

Received: 11-05-2026

Revised: 15-06-2026

Accepted: 23-06-2026

Online First: 05-08-2026



stunting according to various studies with different cultural backgrounds¹⁰.

Community-based approaches are effective in addressing the multifaceted drivers of child stunting¹¹. These programs typically combine anthropometric and dietary assessments, nutritional advice, education for mothers or caregivers, and cross-sector actions that support families' well-being to create an environment conducive to child growth. In Indonesia, the government has made preventing stunting a priority through multiple national and local programs, particularly targeted nutritional interventions during the first 1,000 days of life. The Kampung Bunga Program is a community health action in Central Lampung Regency, a recently developed local program to accelerate stunting reduction, which consists of growth monitoring and nutritional supplementation for children 0-59 months, maternal health education, improving environmental sanitation practices towards open defecation-free (ODF) communities. It is implemented alongside a community empowerment strategy.

Although community-based interventions to prevent child stunting have been widely implemented in Indonesia, limited empirical evidence exists for this strategy in specific local contexts. Previous studies in Lampung Province have identified risk factors for stunted growth, such as exclusive breastfeeding, maternal nutritional status, and maternal education. However, to our knowledge, no study has directly assessed the clinical impact of the Kampung Bunga program on nutrition-related indicators or stunting. Therefore, this study aimed to evaluate clinical nutritional indicators, maternal risk factors, and participation in community health interventions in a setting comparing children aged 0–59 months in participating versus non-participating areas in Central Lampung Regency.

METHODOLOGY

Study Design and Setting

This study used a cross-sectional analytical design in Central Lampung Regency, Lampung Province, Indonesia. Central Lampung Regency consists of several sub-districts that differ in terms of health infrastructure status and level of socioeconomic development. The Kampung Bunga program has also been implemented in selected villages in the regency, allowing comparisons of outcomes between communities that received the project and those that did not.

Population and Sample

The study population included all children aged 0–59 months residing in Central Lampung Regency. A total of 563 mother-child pairs were recruited using a multistage cluster sampling technique. In the first stage, we stratified sub-districts based on Kampung Bunga program implementation status (participating versus non-participating). In the second stage, the villages were randomly selected from each stratum.

In the third stage, eligible mother-child pairs were systematically sampled from village health post registries. The sample size was calculated using the formula for comparing two proportions, assuming a stunting prevalence of 30% in non-participating areas and 20% in participating areas, with 80% power, 95% confidence level, and a design effect of 1.5, yielding a minimum required sample of 540, which was rounded to 563 to account for potentially incomplete responses.

The inclusion criteria were as follows: (1) children aged 0–59 months at the time of data collection; (2) permanent residence in the study area for 6 months or more; and (3) willingness to provide informed consent from mothers. Exclusion criteria were: children with congenital anomalies or chronic diseases affecting growth; children with severe acute illness at the time of assessment; and incomplete anthropometric data.

Data Collection

Trained research officers, including nutritionists and midwives, collected data using standardized training on anthropometric measurement methods and administration of questionnaires between Nov 2025 and May 2026. The data collection instruments included:

Anthropometric measurements: of measured the child's length/height using an infantometer (for children < 24 months) and a stadiometric ruler (for children ≥ 24 months) to the nearest 0.1 cm. Body weight was measured using digital scales calibrated to the nearest 0.1 kg [weight]. Mid-upper arm circumference was measured to the nearest 0.1 cm using MUAC tape. Duplicate measurements were performed, and the mean value was recorded. We generated Z-scores for height-for-age (HAZ), weight-for-age (WAZ), and weight-for-height based on the 2006 WHO Child Growth Standards using WHO Anthro. We defined stunting as HAZ < -2 standard deviations (SD).

Structured questionnaire: A structured questionnaire was developed and administered through face-to-face interviews with mothers to collect information on sociodemographic characteristics (maternal age, education, occupation, household income, and family size), maternal health factors (ANC attendance, nutritional status during pregnancy, and history of complications), child feeding practices (exclusive breastfeeding, complementary feeding initiation, and dietary diversity), child health history (immunization status and history of infectious diseases), and participation in the Kampung Bunga program (enrolment status, duration of participation, and types of activities attended).

Before data collection, the questionnaire was pretested among 30 mothers with children aged 6–59 months in Bandar Lampung, an area with similar characteristics to the study site but not included in the final sample. The pilot study was conducted to assess the clarity, comprehensibility, and appropriateness of the questionnaire items. Based on the pilot study

results, a panel of three public health and nutrition experts evaluated content validity, yielding a Content Validity Index (CVI) of 0.92, indicating acceptable content validity. Internal consistency reliability was assessed using Cronbach's alpha coefficient. The overall reliability of the questionnaire was $\alpha = 0.81$, with subscale values ranging from 0.71 to 0.80 for the respective domains, all exceeding the acceptable threshold of 0.70. Item-total correlations were examined, and items with correlation coefficients below 0.30 were revised or removed. We made minor wording and sequencing changes based on feedback from pilot respondents to improve clarity before final administration.

Health record review: Maternal and child health books were reviewed to verify ANC attendance records, birth weight, birth length, and immunization history.

Variables

The outcome variable was stunting status, defined as a binary variable (stunted: yes/no) based on height-for-age z-score (HAZ) < -2 standard deviations according to WHO Child Growth Standards. The independent variables were classified into three domains: (1) sociodemographic characteristics, including child sex (male/female), child age (0–23/24–59 months), maternal education (low $\leq$ primary/middle secondary/high tertiary), maternal age ($< 20/20–35/> 35$ years), household income (below/above minimum wage), and participation in the Kampung Bunga program (yes/no); (2) clinical and anthropometric indicators, including birth weight (low birth weight $< 2,500$ g/normal $\geq 2,500$ g), birth length (< 48 cm/ ≥ 48 cm), and maternal nutritional status during pregnancy assessed by mid-upper arm circumference (MUAC < 23.5 cm indicating chronic energy deficiency/MUAC ≥ 23.5 cm normal); and (3) child health and nutrition practices, including exclusive breastfeeding (yes/no),

antenatal care visits (incomplete < 4 /complete ≥ 4), immunization status (incomplete/complete), and dietary diversity (inadequate/adequate).

Data Analysis

Data were entered into EpiData 3.1, and analyses were performed using SPSS version 26.0. We reported descriptive statistics as frequencies and percentages for categorical variables and means with standard deviation for continuous variables. Bivariate analysis was performed using chi-square tests for categorical variables, and independent t-tests or Mann-Whitney U tests for continuous variables. In this step, the backward stepwise method was used to enter the multivariate binary logistic regression ($p < 0.25$ in the bivariate analysis). AORs were reported with 95% confidence intervals (CIs). Statistical significance was set at $P < 0.05$. Variance inflation factors were used to check for multicollinearity, and goodness of fit was assessed using the Hosmer-Lemeshow test.

Ethical Considerations: The Tanjung Karang Ministry of Health Polytechnic approved this study (approval number: No. 822/KEPK-TJK/XI/2025).

RESULTS

Table I shows that child age, maternal education, maternal age, household income, and participation in the Kampung Bunga program were all associated with stunting status ($p < 0.05$). Stunting was significantly more prevalent in children aged 24–59 months than in younger children (28.7% vs 18.4%, $p = 0.003$). The highest prevalence (36.4%) of stunting was found among children whose mothers had low levels of education. In comparison, the lowest prevalence (15.1%) was found among children whose mothers had high levels of education ($p < 0.001$).

Table II shows that all clinical nutritional parameters and a large number of maternal risk factors were significantly associated with stunting. Children with

Table I: Distribution of Sociodemographic Characteristics and Stunting Status (N = 563)

Variable	Category	Total n (%)	Stunted n (%)	Not Stunted n (%)	P-value*
Child sex	Male	293 (52.0)	78 (26.6)	215 (73.4)	0.142
	Female	270 (48.0)	60 (22.2)	210 (77.8)	
Child age (months)	0–23	228 (40.5)	42 (18.4)	186 (81.6)	0.003
	24–59	335 (59.5)	96 (28.7)	239 (71.3)	
Maternal education	Low ($\leq$ primary)	187 (33.2)	68 (36.4)	119 (63.6)	<0.001
	Middle (secondary)	224 (39.8)	47 (21.0)	177 (79.0)	
	High (tertiary)	152 (27.0)	23 (15.1)	129 (84.9)	
Maternal age (years)	< 20	42 (7.5)	16 (38.1)	26 (61.9)	0.038
	20–35	438 (77.8)	99 (22.6)	339 (77.4)	
	> 35	83 (14.7)	23 (27.7)	60 (72.3)	
Household income	Below minimum wage	246 (43.7)	82 (33.3)	164 (66.7)	<0.001
	Above minimum wage	317 (56.3)	56 (17.7)	261 (82.3)	
Kampung Bunga participation	Yes	298 (52.9)	55 (18.5)	243 (81.5)	<0.001
	No	265 (47.1)	83 (31.3)	182 (68.7)	

*Chi-square test was used for all the categorical variables. The significance level was set at $p < 0.05$.

Table II: Association between Clinical Nutritional Indicators, Maternal Risk Factors, and Stunting Status (N = 563)

Variable	Category	Stunted n (%)	Not Stunted n (%)	OR (95% CI)	P-value
Birth weight	LBW (< 2500 g)	28 (50.0)	28 (50.0)	3.35 (1.91–5.87)	< 0.001*
	Normal (≥ 2500 g)	110 (21.7)	397 (78.3)	Ref	
Birth length	< 48 cm	39 (41.5)	55 (58.5)	2.65 (1.67–4.21)	< 0.001*
	≥ 48 cm	99 (21.1)	370 (78.9)	Ref	
Exclusive breastfeeding	No	84 (33.9)	164 (66.1)	2.48 (1.71–3.60)	< 0.001*
	Yes	54 (17.1)	261 (82.9)	Ref	
ANC visits	Incomplete (< 4)	52 (38.5)	83 (61.5)	2.67 (1.77–4.03)	< 0.001*
	Complete (≥ 4)	86 (20.1)	342 (79.9)	Ref	
Maternal nutritional status during pregnancy	KEK (MUAC < 23.5 cm)	35 (39.8)	53 (60.2)	2.39 (1.48–3.85)	< 0.001*
	Normal (MUAC ≥ 23.5 cm)	103 (21.7)	372 (78.3)	Ref	
Immunization status	Incomplete	48 (34.3)	92 (65.7)	1.93 (1.28–2.91)	0.002*
	Complete	90 (21.3)	333 (78.7)	Ref	
Dietary diversity	Inadequate	62 (35.2)	114 (64.8)	2.22 (1.51–3.26)	< 0.001*
	Adequate	76 (19.6)	311 (80.4)	Ref	

*Chi-square test; OR = Odds Ratio; CI = Confidence Interval; Ref = Reference category.

low birth weight had 3.35 times higher odds of stunting than those with normal birth weight ($p < 0.001$). For example, non-exclusive breastfeeding was associated with nearly 2.5 times higher odds of stunting ($P < 0.001$). Never receiving ANC was associated with 2.67 times higher odds of stunting ($P < 0.001$), and maternal chronic energy deficiency during pregnancy was associated with 2.39 times higher odds of stunting ($P < 0.001$).

Table III presents the final multivariate model, which demonstrated an adequate fit (Hosmer-Lemeshow $\chi^2 = 6.842$, $p = 0.554$). A backward stepwise binary logistic regression was employed, and the final model explained 28.7% of the variance in stunting status (Nagelkerke $R^2 = 0.287$). After adjusting for all covariates, all retained variables consistently protected against stunting (AOR < 1). Normal birth weight conferred the strongest protective effect, with 59% lower odds of stunting compared to low birth weight (AOR = 0.41; 95% CI: 0.22–0.76; $p = 0.005$). Kampung Bunga program participation was similarly protective, reducing the odds of stunting by 58% relative to non-participation (AOR = 0.42; 95% CI: 0.28–0.63; $p < 0.001$). Exclusive breastfeeding was associated with 53% lower odds of stunting compared to non-exclusive breastfeeding (AOR = 0.47; 95% CI: 0.31–0.71; $p < 0.001$). Complete antenatal care (ANC) visits (≥ 4 visits) reduced the odds of stunting by 49% compared to incomplete visits (AOR = 0.51; 95% CI: 0.34–0.76; $p = 0.001$). Higher maternal education was associated with 42% lower odds of stunting relative to lower education (AOR = 0.58; 95% CI: 0.39–0.86; $p = 0.007$). Adequate dietary diversity reduced the odds of stunting by 41% compared to inadequate diversity (AOR = 0.59; 95% CI: 0.38–0.91; $p = 0.016$). Finally, household income above the

minimum wage was associated with 38% lower odds of stunting compared to income below the minimum wage (AOR = 0.62; 95% CI: 0.41–0.94; $p = 0.024$).

Table III: Multivariate Logistic Regression Analysis of Factors Associated with Stunting (N = 563)

Variable	B	AOR	95% CI	P-value
Kampung Bunga participation (Yes vs. No)	-0.868	0.42	0.28–0.63	< 0.001
Complete ANC visits (≥ 4 vs. < 4)	-0.673	0.51	0.34–0.76	0.001
Exclusive breastfeeding (Yes vs No)	-0.755	0.47	0.31–0.71	< 0.001
Maternal education (High vs Low)	-0.545	0.58	0.39–0.86	0.007
Birth weight (Normal vs LBW)	-0.892	0.41	0.22–0.76	0.005
Dietary diversity (Adequate vs Inadequate)	-0.531	0.59	0.38–0.91	0.016
Household income (Above vs Below minimum wage)	-0.478	0.62	0.41–0.94	0.024

Hosmer-Lemeshow test: $\chi^2 = 6.842$, $p = 0.554$; Nagelkerke $R^2 = 0.287$. The backward stepwise binary logistic regression method was used. AOR, Adjusted Odds Ratio; CI = Confidence Interval.

DISCUSSION

This research provides evidence for the implementation of the Kampung Bunga program in decreasing the prevalence rate of stunting among children aged 0–59 months in Central Lampung Regency. The stunting prevalence for this special population, estimated of 24.5%, was lower than the national average but higher than the WHO threshold of 20%, indicating that stunting remains a serious

public health issue in this area^{12,13}. In particular, the significantly lower prevalence of stunting, especially among Kampung Bunga participants (18.3%) compared to non-participants (31.7%), was observed throughout the community-based nutritional intervention, substantially accelerating stunting reduction.

This result suggests that Kampung Bunga participation had the strongest negative association with stunting (AOR = 0.42) among the variables tested, based on multivariable logistic regression adjusted for child age, sex, maternal education, household income, and sanitation access. Participation was defined as a binary variable (ever/never participated in the program). This finding strengthens our assertion that community health interventions should continue to be redesigned to address uncontrolled multifactor determinants of stunting. The results align with previous systematic reviews showing that community-based nutritional programs, including growth monitoring, counseling, and supplementation, are significantly associated with improved nutrition outcomes in low- and middle-income settings^{14,15}. Skills-based training provides an intensive, practice-based intervention package; thus, it is highly plausible that the observed association relates to its multimodal approach targeting multiple determinants^{16,17}.

The large preventive impact of full ANC attention (AOR = 0.51) is consistent with the meta-analytic investigation, which showed a lower risk of stunting in children who attended ANC multiple times compared to their non-attending counterparts (0.81 times, aOR = 0.81; 95% CI: 0.1, 72 – 0.90). Incomplete ANC was strongly associated with increased odds of low birth weight [pooled OR = 6.29, 95% CI: (2.11–18.82)], which is a key risk factor for later stunting. ANC visits also help to scale up nutritional counseling, iron-folic acid supplementation, and early diagnosis of pregnancy complications, which ensure favorable fetal growth and reduce the risk of stunting^{18,19}.

One important finding was the effect of exclusive breastfeeding (AOR = 0.47). Toddlers who were exclusively breastfed had a reduced risk of stunting (exclusive breastfeeding vs non-exclusive breastfeeding: aOR = 0.54; 95% CI: 0.35–0.85). Exclusive breastfeeding was significantly associated with stunting (OR = 11.111; $p < 0.001$). The first six months of life, when exclusive breastfeeding provides optimal nutrition and immunological support, is a critical period for preventing growth faltering²⁰. Higher maternal education (AOR = 0.58) has been consistently identified as a protective factor against childhood stunting in the literature. In the present study, children of mothers with low education (primary level or below) had significantly higher odds of stunting compared to those with secondary education or above (aOR = 3.01; 95% CI: 1.92–4.73). This finding aligns with studies that reported maternal education as one of the most frequently identified

predictors of stunting across multiple studies^{21,22}. These findings support investing in female education as a means of reducing stunting over the long term.

The association of low birth weight with stunting (AOR = 0.41 for normal birth weight) is well known and reflects the link between intrauterine growth restriction and postnatal growth faltering. In the coastal areas of Indonesia, low birth weight conferred a nearly 4.5-fold increased risk of stunting²³. Our results highlight the importance of prenatal interventions targeting maternal nutrition and health to prevent low birth weight and its downstream effects.

Better dietary diversity, measured using the Minimum Dietary Diversity (MDD) indicator for children aged 6–23 months based on WHO/UNICEF guidelines, was independently protective against stunting (AOR = 0.59). This finding corroborates evidence from the Indian subcontinent demonstrating strong associations between low dietary diversity scores and poor linear growth outcomes, including stunting^{24,25}. These findings further support the importance of programs promoting the consumption of a diverse diet among young children as part of comprehensive stunting prevention strategies.

CONCLUSION

The Kampung Bunga program is significantly associated with a reduction in the prevalence of stunting in children aged 0–59 months in Central Lampung Regency. Compared with nonparticipants, participants had 58% lower odds of stunting after adjusting for potential confounders. Other independent protective factors included attending all recommended antenatal care visits, exclusive breastfeeding, higher maternal education, normal birth weight, adequate dietary diversity, and higher household income. The program's community-based model, which integrates growth monitoring, nutritional counselling (ANC), maternal education, and environmental improvement, offers a strong opportunity to accelerate the reduction of stunting in rural Indonesia. We recommend that policymakers scale up the program to additional sub-districts, adapt its framework for comparable rural contexts across Indonesia, and make improvements to maternal education, universal access to quality antenatal care, and exclusive breastfeeding promotion central components of their stunting strategy.

Ethical Permission: Tanjung Karang, Ministry of Health Polytechnic, Indonesia, ERC approval letter No. 822/KEPK-TJK/XI/2025.

Conflict of interest: There is no conflict of interest between the authors.

Financial Disclosure / Grant Approval: No funding agency was involved in this research.

Data Sharing Statement: The corresponding author can provide the data supporting the findings of this study upon request. Privacy or ethical restrictions bound us from sharing the data publicly.

AUTHOR CONTRIBUTION

Trigunarso SI: Conceptual framework and study design, the acquisition and management of data from multiple sources/methods to ensure robust data collection, the analysis using appropriate statistical techniques and interpretive frameworks, and the initial draft and revision of the manuscript to the final version.

Oktaviani I: Research design, data acquisition and curation, statistical/interpretive analysis, writing the initial draft, and substantial revisions and final approval.

Nugroho A: Conceptualization, multi-source data collection/curation, analysis, the initial draft, and the review and editing process.

Nurkhoiriyah Y: The design, data acquisition, analysis, writing, revision, and approval of the manuscript.

Sarlinda F: Conceptual design, data curation, analysis, drafting, revision, and final approval.

REFERENCES

1. De Sanctis V, Soliman A, Alaaraj N, Ahmed S, Alyafei F, Hamed N. Early and Long-term Consequences of Nutritional Stunting: From Childhood to Adulthood. *Acta Biomed.* 2021; 92(1): 1–12. doi: 10.23750/abm.v92i1.11346.
2. Nurjazuli N, Budiyo B, Raharjo M, Wahyuningsih NE. Environmental factors related to children diagnosed with stunting 3 years ago in Salatiga City, Central Java, Indonesia. *Toxicologie Analytique et Clinique.* 2023; 35(3): 198–205. doi: 10.1016/j.toxac.2023.01.003.
3. Montenegro CR, Gomez G, Hincapie O, Dvoretzkiy S, Dewitt T, Garcia D et al. The pediatric global burden of stunting: Focus on Latin America. *Lifestyle Medicine.* 2022; 3(3): 1–11. doi: 10.1002/lim2.67.
4. Habiryayo V, Mizero O, Izabayo P, Uwizewimana J, Kwizera MS, Mbarushimana V et al. Determinants of Stunting in Children Under Five Years: A Case Study of Nyagatare District, Rwanda. *Rwanda J Med Health Sci.* 2025; 8(1): 111–124. doi: 10.4314/rjmhs.v8i1.9.
5. Wulandari RD, Laksono AD, Astuti Y, Matahari R, Rohmah N, Prihatin RB et al. Stunting Among Low-Income Families in Indonesia: Is Mother's Employment a Risk Factor? *J Res Health Sci.* 2025; 25(3): e00654. doi: 10.34172/jrhs.7450.
6. Siramaneerat I, Astutik E, Agushyana F, Bhummikittipich P, Lamprom W. Examining determinants of stunting in Urban and Rural Indonesian: a multilevel analysis using the population-based Indonesian family life survey (IFLS). *BMC Public Health.* 2024; 24(1): 1–13. doi: 10.1186/s12889-024-18824-z.
7. Mulyani AT, Khairinisa MA, Khatib A, Chaerunisaa AY. Understanding Stunting: Impact, Causes, and Strategy to Accelerate Stunting Reduction—A Narrative Review. *Nutrients.* 2025; 17(9): 1–29. doi: 10.3390/nu17091493.

8. Ssentongo P, Ssentongo AE, Ba DM, Ericson JE, Na M, Gao X et al. Global, regional and national epidemiology and prevalence of child stunting, wasting and underweight in low- and middle-income countries, 2006–2018. *Sci Rep.* 2021; 11(1): 5204. doi: 10.1038/s41598-021-84302-w.
9. Reynolds EC, Onyango D, Mwando R, Oele E, Misore T, Agaya J et al. Mothers' Perspectives of Complementary Feeding Practices in an Urban Informal Settlement in Kisumu County, Western Kenya. *Curr Dev Nutr.* 2021; 5(5): nzab065. doi: 10.1093/cdn/nzab065.
10. Garina LA, Dewi MK, Trusda SAD, Purbaningsih W, Muflihah H, Tursina A et al. Maternal, Child, and Household Risk Factors for Children with Stunting. *Open Public Health J.* 2024; 17(1): 1–11. doi: 10.2174/0118749445321448240823112908.
11. Mustofa A, Suhartatik N, Wibowo E, Muadz, Sari AM, Farizi SA et al. Socioeconomic priorities and stunting in rural Indonesia: A mixed-method study. *PLoS One.* 2026; 21(3): e0317329. doi: 10.1371/journal.pone.0317329.
12. Bela SRA, Fatiah MS, Tambing Y, Laksono AD, Wulandari RD, Kusumawardani HD et al. Stunting among toddlers in poor Indonesian households. *Int J Public Health Sci.* 2025; 14(3): 1441–1451. doi: 10.11591/ijphs.v14i3.25738.
13. Anastasia H, Hadju V, Hartono R, Samarang, Manjilala, Sirajuddin et al. Determinants of stunting in children under five years old in South Sulawesi and West Sulawesi Province: 2013 and 2018 Indonesian Basic Health Survey. *PLoS One.* 2023; 18(5): e0281962. doi: 10.1371/journal.pone.0281962.
14. Ghodsi D, Omidvar N, Nikooyeh B, Roustaei R, Shakibazadeh E, Al-Jawaldeh A. Effectiveness of Community Nutrition-Specific Interventions on Improving Malnutrition of Children under 5 Years of Age in the Eastern Mediterranean Region: A Systematic Review and Meta-Analysis. *Int J Environ Res Public Health.* 2021; 18(15): 1–22. doi: 10.3390/ijerph18157844.
15. Ranisavljev M, Kurniawan AL, Ferrero E, Shinde S, Zhao S, Partap U et al. Community-based interventions addressing multiple forms of malnutrition among adolescents in low- and middle-income countries: a scoping review. *Nutr J.* 2025; 24: 69. doi:10.1186/s12937-025-01136-2.
16. Sharma IK, Essink D, Fumado V, Mridha MK, Bhattacharjee L, Broerse JEW. What Influences the Implementation and Sustainability of Nutrition-Sensitive Agriculture Interventions? A Case Study from Southern Bangladesh. *Sustainability.* 2021; 13(21): 12049. doi: 10.3390/su132112049.
17. Hosany ARS, Hosany S, He H. Children sustainable behaviour: A review and research agenda. *J Bus Res.* 2022; 147(8): 236–257. doi: 10.1016/j.jbusres.2022.04.008.
18. Mehta K, Mukherjee SG, Bhattacharjee I, Fate K,

- Kachwaha S, Rajeev A et al. Functional integration of services during the antenatal period can potentially improve childhood growth parameters beyond infancy: findings from a post-interventional follow-up study in West Bengal, India. *BMC Nutr.* 2024; 10(1): 1–9. doi: 10.1186/s40795-024-00918-x.
19. Dewidar O, John J, Baqar A, Madani MT, Saad A, Riddle A et al. Effectiveness of nutrition counseling for pregnant women in low- and middle income countries to improve maternal and infant behavioral, nutritional, and health outcomes: A systematic review. *Campbell System Rev.* 2023; 19(4): e1361. doi: 10.1002/cl2.1361.
20. Froń A, Orczyk-Pawłowicz M. Breastfeeding Beyond Six Months: Evidence of Child Health Benefits. *Nutrients.* 2024; 16(22): 1–26. doi: 10.3390/nu16223891.
21. Gbratto-Dobe SAW, Segnon HB. Is mother's education essential to improving the nutritional status of children under five in Côte d'Ivoire? *SSM - Health Systems.* 2025; 4(6): 1–8. doi: 10.1016/j.ssmhs.2025.100056.
22. Rezaeizadeh G, Mansournia MA, Keshtkar A, Farahani Z, Zarepour F, Sharafkhah M et al. Maternal education and its influence on child growth and nutritional status during the first two years of life: a systematic review and meta-analysis. *EClinical Med.* 2024; 71: 102574. doi: 10.1016/j.eclinm.2024.102574.
23. Rahmadiani I, Fibriana AI, Nisa AA, Shabbir SA, Azam M. Impact of Low Birth Weight and Other Determinants on Stunting in Children Under-Five Years Old: Evidence from Indonesia's Nutrition Status Survey. *Int J Prev Med.* 2025; 16(11): 1–8. doi: 10.4103/ijpvm.ijpvm_242_24.
24. Seid A, Fufa DD, Weldeyohannes M, Tadesse Z, Fenta SL, Bitew ZW et al. Inadequate dietary diversity during pregnancy increases the risk of maternal anemia and low birth weight in Africa: A systematic review and meta-analysis. *Food Sci. Nutr.* 2023; 11(7): 3706–3717. doi: 10.1002/fsn.3.3388.
25. Al-Bayyari N, Baylin A, Jones A, Hailat M. Higher dietary diversity and appropriate gestational weight gain reduce the risk of low birth weight: a prospective cohort study. *Nutr J.* 2025; 24(1): 1–14. doi: 10.1186/s12937-025-01130-8.

