

# Quality of Life among Adults Undergoing Sleeve Gastrectomy and Lifestyle Modifications: Comparative Analytical Cross-Sectional Study

Noman Dilshad<sup>1</sup>, Pamela Marshall<sup>2</sup>, Shakirullah<sup>3\*</sup>

## ABSTRACT

**OBJECTIVE:** To assess the health-related Quality of life of adult patients undergoing sleeve gastrectomy with lifestyle modifications.

**METHODOLOGY:** A Comparative, Analytical, cross-sectional design was employed at Ziauddin and Civil Hospitals, Karachi, from June to November 2022. A total of 102 participants were recruited via consecutive sampling, with 51 in the sleeve gastrectomy group and 51 in the lifestyle modification group. Inclusion criteria included adults aged 18 years or older who had undergone sleeve gastrectomy or followed lifestyle modification for at least 3 months. Exclusion criteria were patients with terminal illnesses (e.g., cancer, HIV/AIDS) and those with diagnosed psychiatric disorders. Data were collected using the RAND SF-36 questionnaire. Statistical analysis was performed via SPSS version 26.

**RESULTS:** Among females, the mean score was 53.79 (SD  $\pm$  3.3) with sleeve gastrectomy and a mean score of 37.46 (SD  $\pm$  7.7) with lifestyle modification (p-value < 0.001). Furthermore, in males, the mean score was 53.67  $\pm$  4.2 with sleeve gastrectomy and a mean score of 40.09  $\pm$  9.5 with lifestyle modification (p-value < 0.001).

**CONCLUSION:** The findings of the study found that 37 (37.3%) of the participants had a very good quality of life, while 8% reported poor Quality of life with both procedures. When comparing the quality-of-life score with the procedure of sleeve gastrectomy and lifestyle modification, it was reported as significant with a p-value of less than 0.05.

**KEYWORDS:** Sleeve Gastrectomy, Lifestyle Modifications, Quality of Life, Surgery, Patients, Gastrectomy Procedure.

## INTRODUCTION

The globally emerging epidemic of non-communicable diseases (NCDs) is creating major health challenges and is threatening to overwhelm the healthcare delivery system. Non-communicable diseases NCDs are becoming a crisis and are being recognized as a leading cause of mortality and morbidity worldwide<sup>1</sup>. Among the different causes of NCDs, obesity or overweight (increased body mass index) is considered a major cause and a leading health problem that increases the probability of getting NCDs and other health problems<sup>2</sup>. Multiple factors contribute to obesity, including dietary patterns, physical inactivity, genetic predisposition, social roles, and others, leading to a higher proportion of body fat than is required<sup>3</sup>.

Obesity is accountable for 2.8 million deaths annually throughout the world, and approximately 35.8 million (2.3%) of the world population is obese<sup>4</sup>. However, the

risk of obesity is intensifying in all countries and among all age groups, genders, and income groups<sup>5</sup>. A study conducted in Chile reported that high BMI  $\geq$  25 kg/m<sup>2</sup> accounted for 31.6% of NCD deaths, with global estimates of 4.7 million NCDs and a projected annual rise<sup>6</sup>. Another comprehensive review reported that over 70% of NCDs have an association with obesity, which highlights the pervasive impact of excess weight on the risk of chronic diseases<sup>7</sup>. It is disproportionately affecting middle-income countries (LMICs), posing a double burden on these countries for the diseases since the last decade<sup>8</sup>. Research conducted in Pakistan suggests there is a rising trend of obesity among the community population. The findings report an obesity prevalence of 22.0%<sup>9</sup>. Moreover, the literature documents that obesity alters an individual's biological, physiological, psychological, behavioral, and social processes. The process may result in restriction or lack of ability to perform an activity with flexibility or within the range considered normal<sup>10,11</sup>. In addition, this negatively affects functional capability and social habits and is considered a main predictor of poor Quality of life<sup>12</sup>. Globally, various treatment methods and techniques have been adopted to reduce obesity or its consequences, such as increased BMI, including exercise, medications, low-fat and low-carbohydrate diets, reduced calorie intake, surgery, and lifestyle

<sup>1</sup>Assistant Professor/ Principal, Jesus & Mary College of Nursing, Karachi, Pakistan

<sup>2</sup>Dean, Ziauddin University Faculty of Nursing and Midwifery

<sup>3</sup>Ph.D. Scholar, Ziauddin University Faculty of Nursing and Midwifery

**Correspondence:** shakirullah.19940@zu.edu.pk

doi: 10.22442/jlumhs.2026.01380

Received: 27-08-2025

Revised: 02-03-2026

Accepted: 09-03-2026

Online First: 06-05-2026



modifications<sup>13,14</sup>. Among these techniques, sleeve gastrectomy surgery has emerged as the most desired and effective technique to counter obesity and its associated symptoms. The surgery is mainly recommended for individuals with a BMI of 30-34.9 kg/m<sup>2</sup> or for those who fail to adopt the above-mentioned treatment option<sup>15</sup>. On the other hand, scientific findings also suggest that individuals with obesity prefer lifestyle modification methods to reduce their increased BMI and related complications<sup>16</sup>.

Despite the evidence that has highlighted the effect of obesity on NCDs, limited comparative studies exist within the Pakistani context evaluating health-related quality of life among patients undergoing lifestyle modification and sleeve gastrectomy. Given the limited evidence from low- and middle-income countries, this study aims to address this gap by comparing QoL outcomes among adults undergoing sleeve gastrectomy or lifestyle modification at a tertiary care hospital in Karachi.

The current study aims to compare the Quality of life (QOL) of adult patients undergoing sleeve gastrectomy versus lifestyle modification. To determine the possible determinants of the quality of life of adult patients having sleeve gastrectomy in comparison to lifestyle modification.

## METHODOLOGY

A Comparative Analytical Cross-Sectional design was employed in the current study. The study was conducted at Ziauddin University Hospital (ZUH), Karachi, a private and public tertiary care teaching hospital, and Civil Hospital Karachi. The duration of the study was from June to November 2022, after the approval of all ethical and research committees. The target population was patients who had undergone sleeve gastrectomy or adopted lifestyle modifications to control increasing weight. Only participants who had adopted a nutritional method for at least of month or had undergone sleeve gastrectomy were included. A consecutive sampling technique was used to recruit the study participants. The sample size was calculated using the Online Software OpenEpi based on a census of 2 months. The sample size was calculated with a 95% confidence interval and a 5% margin of error of 5%. The calculated sample size was 102, equally divided into two groups: 51 in the sleeve gastrectomy group and 51 in the lifestyle modification group. The following inclusion and exclusion criteria were used to include participants in the current study. Patients aged 18 years and above, with no upper age limit, regardless of gender, who underwent sleeve gastrectomy. Those adopting any lifestyle modification were included.

Participants with any terminal illness, such as cancer, HIV (AIDS), or others, were excluded. Participants suffering from any mental or psychiatric conditions were excluded.

The current study used a valid and reliable data collection tool. The first component of the

questionnaire included demographics and questions about co-morbidities. The second component consisted of the RAND-developed Short Form Health Survey (SF-36). The tool is structured, organized, and coherent. It is a Likert scale that measures quality of life across eight domains of health.

This procedure began after obtaining permission to use the instrument and approval from all concerned committees and the study setting. The selected participants were approached, and the date and time for completing the tool were decided. The study's purpose and objectives were then shared with potential participants. Eligible participants were identified and invited to participate voluntarily by signing the informed consent form. Afterwards, the data were collected using a study tool provided to the participants.

The data were analyzed using SPSS 26, and both descriptive and inferential analyses were performed. Initially, the data were checked for normality through the Shapiro–Wilk test. Normally distributed data were analyzed using parametric tests. In descriptive statistics, means  $\pm$  standard deviations were calculated for continuous variables, and frequencies and proportions were reported for categorical variables. In the inferential analysis, the difference in the mean score for continuous QoL variables was assessed using a t-test for two independent samples, and the difference in proportions for categorical variables was assessed using a Chi-square test of independence. The analysis was performed at a 95% confidence level (level of significance of 0.05), and a P-value of less than 0.05 was considered statistically significant.

## Ethical Consideration

The study obtained approval from the university's Ethical Review Committee (Reference No. 4960222NDNUR) and permission to use the tools from the developers. Permission to use the study setting was obtained from the hospitals, and a consultant was approached to recruit the study participants.

## RESULTS

**Table I: Socio-demographic characteristics of study participants**

| Demographic Variable<br>n=102 (100%) | Frequency         | Percent |
|--------------------------------------|-------------------|---------|
| <b>Gender</b>                        |                   |         |
| Male                                 | 38                | 37.25%  |
| Female                               | 64                | 62.75%  |
| <b>Study Setting</b>                 |                   |         |
| Ziauddin Hospital                    | 20                | 19.61%  |
| Civil Hospital                       | 82                | 80.39%  |
| <b>Mean</b>                          |                   |         |
| Age (years)                          | 35.84 $\pm$ 5.7   |         |
| Weight (per KG)                      | 143.03 $\pm$ 13.3 |         |
| Height (feet)                        | 5.81 $\pm$ 0.25   |         |

| <b>Employment Setting</b>            |    |        |
|--------------------------------------|----|--------|
| Self-Employed                        | 30 | 29.41% |
| Part Timer                           | 14 | 13.73% |
| A homemaker                          | 11 | 10.78% |
| Student                              | 13 | 12.75% |
| Military                             | 8  | 7.84%  |
| Retired                              | 1  | 0.98%  |
| Other                                | 25 | 24.51% |
| <b>Informal Education</b>            |    |        |
| Madrasa                              | 12 | 11.76% |
| Urdu Reading only                    | 9  | 8.82%  |
| Certificate received for the diploma | 2  | 1.96%  |
| Diploma without a certificate        | 38 | 37.26% |
| Degree                               | 41 | 40.20% |
| <b>Language</b>                      |    |        |
| Urdu                                 | 46 | 45.10% |
| Sindhi                               | 16 | 15.69% |
| Balochi                              | 7  | 6.86%  |
| Pashto                               | 11 | 10.78% |
| Punjabi                              | 18 | 17.65% |
| Siraiki                              | 4  | 3.92%  |
| <b>Family Status</b>                 |    |        |
| Nuclear                              | 56 | 54.90% |
| Extended                             | 41 | 40.20% |
| Single                               | 5  | 4.90%  |
| <b>Marital Status</b>                |    |        |
| Single                               | 7  | 6.86%  |
| Married                              | 86 | 84.32% |
| Divorced/Separated                   | 8  | 7.84%  |
| Widowed                              | 1  | 0.98%  |
| <b>Socioeconomic Status</b>          |    |        |
| Lower                                | 11 | 10.78% |
| Lower middle Class                   | 13 | 12.75% |
| Middle Class                         | 69 | 67.65% |
| Upper Class                          | 9  | 8.82%  |

"Percentages were rounded to two decimal places, and minor adjustments were made to ensure totals equal 100%"

#### **Comparisons of Quality-of-Life Score with respect to Sleeve gastrectomy & lifestyle modification**

The quality-of-life scores of participants who underwent the procedure of sleeve gastrectomy and lifestyle modification were analyzed. A p-value of less than 0.05 was considered significant. According to the study results, the average quality-of-life score with sleeve gastrectomy was higher, with a mean of  $53.7 \pm 4$ , than with lifestyle modification,  $38.9 \pm 8.8$ . A higher score shows a better quality of life. The result is shown in **Table II** (P-value <0.001).

**Table II: Comparisons of Quality-of-Life Score with respect to Sleeve gastrectomy & Lifestyle modification**

| Method n=102                         | Quality Of Life | P-value |
|--------------------------------------|-----------------|---------|
|                                      | Mean $\pm$ SD   |         |
| <b>Sleeve gastrectomy (n=51)</b>     | 53.7 $\pm$ 4    | <0.001  |
| <b>Lifestyle Modification (n=51)</b> | 38.9 $\pm$ 8.8  |         |

An Independent T-test has been applied. A P-value  $\leq 0.05$  is considered significant.

**Table III: Comparisons of Quality of Life Domains with respect to sleeve gastrectomy & lifestyle modification**

| Quality of Life                | Sleeve Gas-trectomy n=51 |        | Lifestyle Modifi-cation n=51 |        | P-Value |
|--------------------------------|--------------------------|--------|------------------------------|--------|---------|
| General Health                 |                          |        |                              |        |         |
| Poor                           | 0                        | 0.00%  | 8                            | 15.69% | <0.001  |
| Fair                           | 0                        | 0.00%  | 14                           | 27.45% |         |
| Good                           | 11                       | 21.57% | 24                           | 47.06% |         |
| Very Good                      | 35                       | 68.63% | 3                            | 5.88%  |         |
| Excellent                      | 5                        | 9.80%  | 2                            | 3.92%  |         |
| Limitations of activities      |                          |        |                              |        |         |
| Yes, limited a lot             | 0                        | 0.0%   | 4                            | 7.84%  | 0.041   |
| Limited a little               | 51                       | 100%   | 45                           | 88.24% |         |
| No                             | 0                        | 0.0%   | 2                            | 3.92%  |         |
| Physical health problem        |                          |        |                              |        |         |
| Yes                            | 21                       | 41.18% | 8                            | 15.69% | 0.004   |
| No                             | 30                       | 58.82% | 43                           | 84.31% |         |
| Emotional Health Problem       |                          |        |                              |        |         |
| Yes                            | 0                        | 0.00%  | 12                           | 23.53% | <0.001  |
| No                             | 51                       | 100%   | 39                           | 76.47% |         |
| Social activities interference |                          |        |                              |        |         |
| Most of the Time               | 2                        | 3.92%  | 17                           | 33.33% | <0.001  |
| Some of the Time               | 31                       | 60.79% | 34                           | 66.67% |         |
| A little of the Time           | 17                       | 33.33% | 0                            | 0.00%  |         |
| None of the Time               | 1                        | 1.96%  | 0                            | 0.00%  |         |
| Pain                           |                          |        |                              |        |         |
| Severe                         | 0                        | 0.00%  | 21                           | 41.18% | <0.001  |
| Moderate                       | 7                        | 13.73% | 17                           | 33.33% |         |
| Mild                           | 43                       | 84.31% | 13                           | 25.49% |         |
| Very Mild                      | 1                        | 1.96%  | 0                            | 0.00%  |         |
| Energy and emotional fall      |                          |        |                              |        |         |
| Most of the time               | 0                        | 0.00%  | 3                            | 5.88%  | <0.001  |
| A good bit of the time         | 1                        | 1.96%  | 35                           | 68.63% |         |
| Some of the time               | 39                       | 76.47% | 9                            | 17.65% |         |
| A little of the time           | 11                       | 21.57% | 3                            | 5.88%  |         |
| None of the time               | 0                        | 0.00%  | 1                            | 1.96%  |         |
| Low Mental Health              |                          |        |                              |        |         |
| Mostly True                    | 0                        | 0.00%  | 1                            | 1.96%  | <0.001  |
| Don't Know                     | 15                       | 29.41% | 39                           | 76.47% |         |
| Mostly False                   | 36                       | 70.59% | 10                           | 19.61% |         |
| False                          | 0                        | 0.00%  | 1                            | 1.96%  |         |

The chi-Square test has been applied. A P-value  $\leq 0.05$  is considered significant.

*"Percentages were rounded to two decimal places, and minor adjustments were made to ensure totals equal 100%"*

**Table IV: Comparison of Quality-of-Life score among Demographic Variables with respect to Sleeve gastrectomy and Lifestyle modification**

| Demographic Variables n=102          | Status                 | SG=51<br>LSM=51 | Mean ±<br>std | P-<br>value |
|--------------------------------------|------------------------|-----------------|---------------|-------------|
| Gender                               |                        |                 |               |             |
| Female                               | Sleeve Gastrectomy     | 14              | 53.79 ±3.3    | <0.001      |
|                                      | Lifestyle Modification | 24              | 37.46 ±7.7    |             |
| Male                                 | Sleeve Gastrectomy     | 37              | 53.67 ±4.2    | <0.001      |
|                                      | Lifestyle Modification | 27              | 40.09 ±9.5    |             |
| Hospital Setting                     |                        |                 |               |             |
| Civil Hospital                       | Lifestyle Modification | 20              | 37.92 ±6.8    | <0.001      |
| Ziauddin Hospital                    | Sleeve Gastrectomy     | 51              | 53.70 ±4.0    | <0.001      |
|                                      | Lifestyle Modification | 31              | 39.45 ±9.9    |             |
| Employment Setting                   |                        |                 |               |             |
| Self-Employed                        | Sleeve Gastrectomy     | 12              | 53.69 ±2.3    | <0.001      |
|                                      | Lifestyle Modification | 19              | 37.68 ±8.7    |             |
| Part Timer                           | Sleeve Gastrectomy     | 9               | 54.08 ±3.9    | <0.001      |
|                                      | Lifestyle Modification | 4               | 39.82 ±6.7    |             |
| A homemaker                          | Sleeve gastrectomy     | 6               | 51.47 ±4.8    | 0.077       |
|                                      | Lifestyle Modification | 5               | 39.12 ±14.4   |             |
| Student                              | Sleeve gastrectomy     | 8               | 54.20 ±3.7    | 0.081       |
|                                      | Lifestyle Modification | 5               | 42.83 ±16.5   |             |
| Military                             | Sleeve gastrectomy     | 2               | 54.42 ±8.1    | 0.004       |
|                                      | Lifestyle Modification | 6               | 36.28 ±3.9    |             |
| Retired                              | Lifestyle Modification | 1               | 43.36 ±3.9    | <0.001      |
| Other                                | Sleeve gastrectomy     | 14              | 54.05 ±4.7    |             |
|                                      | Lifestyle Modification | 11              | 39.58 ±4.2    |             |
| Informal Education                   |                        |                 |               |             |
| Madrassa                             | Sleeve gastrectomy     | 5               | 53.10 ±5.7    | 0.001       |
|                                      | Lifestyle Modification | 7               | 38.81 ±4.6    |             |
| Urdu Reading only                    | Sleeve gastrectomy     | 9               | 53.59 ±4.5    |             |
| Certificate received for the diploma | Lifestyle Modification | 2               | 31.86 ±0.0    |             |

|                               |                        |    |             |        |
|-------------------------------|------------------------|----|-------------|--------|
| Diploma without a certificate | Sleeve gastrectomy     | 13 | 52.48 ±4.1  | <0.001 |
|                               | Lifestyle Modification | 25 | 39.82 ±9.5  |        |
| Degree                        | Sleeve gastrectomy     | 24 | 54.54 ±3.3  | <0.001 |
|                               | Lifestyle Modification | 17 | 38.26 ±9.4  |        |
| Language                      |                        |    |             |        |
| Urdu                          | Sleeve gastrectomy     | 29 | 54.01 ±3.6  | <0.001 |
|                               | Lifestyle Modification | 17 | 40.24 ±10.8 |        |
| Sindhi                        | Sleeve gastrectomy     | 5  | 53.45 ±5.5  | <0.001 |
|                               | Lifestyle Modification | 11 | 34.59 ±5.2  |        |
| Balochi                       | Sleeve gastrectomy     | 3  | 49.56 ±4.9  | 0.055  |
|                               | Lifestyle Modification | 4  | 38.50 ±6.3  |        |
| Pashto                        | Sleeve gastrectomy     | 5  | 54.51 ±5.6  | 0.001  |
|                               | Lifestyle Modification | 6  | 37.17 ±6.0  |        |
| Punjabi                       | Sleeve gastrectomy     | 7  | 54.24 ±2.7  | 0.006  |
|                               | Lifestyle Modification | 11 | 41.67 ±10.1 |        |
| Siraiki                       | Sleeve gastrectomy     | 2  | 52.21 ±2.5  | <0.001 |
|                               | Lifestyle Modification | 2  | 40.71 ±3.8  |        |
| Family Status                 |                        |    |             |        |
| Nuclear                       | Sleeve gastrectomy     | 32 | 54.04 ±3.6  | <0.001 |
|                               | Lifestyle Modification | 24 | 39.71 ±10.7 |        |
| Extended                      | Sleeve gastrectomy     | 17 | 53.31 ±4.8  | <0.001 |
|                               | Lifestyle Modification | 24 | 37.79 ±7.1  |        |
| Single                        | Sleeve gastrectomy     | 2  | 51.77 ±1.9  | 0.007  |
|                               | Lifestyle Modification | 3  | 40.41 ±1.8  |        |
| Marital Status                |                        |    |             |        |
| Single                        | Sleeve gastrectomy     | 3  | 56.64 ±1.5  | 0.364  |
|                               | Lifestyle Modification | 4  | 47.12 ±16.0 |        |
| Married                       | Sleeve gastrectomy     | 41 | 53.49 ±4.0  | <0.001 |
|                               | Lifestyle Modification | 45 | 38.03 ±7.9  |        |
| Divorced/ Separated           | Sleeve gastrectomy     | 7  | 53.73 ±4.1  | 0.015  |
|                               | Lifestyle Modification | 1  | 38.94 ±3.3  |        |
| Widowed                       | Lifestyle Modification | 1  | 42.48 ±3.5  |        |
| Socioeconomic Status          |                        |    |             |        |
| Lower                         | Lifestyle Modification | 11 | 37.81 ±5.3  |        |
| Lower middle Class            | Sleeve gastrectomy     | 7  | 53.22 ±4.2  | <0.001 |
|                               | Lifestyle Modification | 6  | 36.43 ±6.0  |        |
| Middle Class                  | Sleeve gastrectomy     | 39 | 53.85 ±4.0  | <0.001 |
|                               | Lifestyle Modification | 30 | 40.59 ±10.0 |        |
| Upper Class                   | Sleeve gastrectomy     | 5  | 53.27 ±3.9  | 0.001  |
|                               | Lifestyle Modification | 4  | 32.30 ±6.6  |        |

*An Independent T-test has been applied. A P-value  $\leq 0.05$  is considered significant.*

## DISCUSSION

According to the study results, the participants were  $n=102$ , with a mean age of 35.84 years and an SD of  $\pm 5.7$ . These findings were similar to those of a study by Timkova V 2024<sup>17</sup> in 2017, in which the mean age of participants was 44.9 years, with an SD of  $\pm 12.7$ . The mean weight of the participants was 143.03 kg (SD of  $\pm 13.3$ ), and the mean height of the participants was 5.81 feet (SD of  $\pm 0.25$ ). These findings were similar to the results of a study in which the mean weight of the participants was 133.3kg (SD  $\pm 15.3$ )<sup>18</sup>. By gender, the subjects were categorized into two groups: 37.3% of the participants were female, and 62.7% were male. Similarly, these findings were similar to those of a study by Castro Al et al.<sup>19</sup> in 2018, in which 60% of participants were female, and 40% were male. By employment status of the participants, 29.4% of the participants were self-employed, 13% were part-timers, 10.8% were homemakers, 12.7% were students, 7.8% were military, 1% were retired, and 24.5% were from another profession. These findings contradicted the results of a 2018 study by Rafique I et al.<sup>4</sup>, in which the majority (62.8%) were unpaid, followed by 21.3% non-government employees, 12.3% self-employed, and 3.7% government employees. In the current study, regarding educational status, the majority of participants (40.2%) held degrees, followed by 37.3% with a diploma or certificate, 11.8% who had attended madrasa, and 8.85% who could read Urdu only. These findings were similar to those of a study conducted by Akkayaoğlu H 2020<sup>20</sup>, in which the majority (44%) of participants had university-level education, followed by 42% of participants with a high school level of education, and 14% with an elementary level of education. By ethnicity, most participants (45.1%) were Urdu speakers, followed by Sindhi (15.7%), Pashto (17.6%), and other ethnic backgrounds. These findings were similar to those of a study conducted in Malaysia, which included participants from different ethnic backgrounds<sup>5</sup>. Furthermore, the marital status of the participants showed that most participants (84.3%) were married, 7% were single, 8% were divorced, and only 1% were widowed. These findings were similar to those of a study in which most participants (80%) were married, and 20% were unmarried<sup>19</sup>. Likewise, another study showed the same findings: 86.2% of married participants and 13.8% of unmarried participants<sup>5</sup>. According to the study findings, socioeconomic statuses of participants were divided into four categories: 10.8% ( $n=11$ ) belonged to the lower Class, 12.7% ( $n=13$ ) belonged to the lower middle Class, 67.6% ( $n=69$ ) belonged to the middle Class, and the other 8.8% ( $n=9$ ) belonged to the upper Class. (Table I)

### Analysis of Quality of Life, comparison of Quality-of-life score, and Quality of Life with respect to the procedure

The Quality of Life of the participants were analyzed

through the SF-36 Scale. The results of the study showed that the majority (37.3%) of the participants had a very good quality of life, followed by 34.3% with a good quality of life, 6.9% with an excellent quality of life, and 13.7% with a fair quality of life. In comparison, 78% reported poor quality of life, indicating a large gap in the overall level of life satisfaction and well-being of the group investigated. Conversely, these findings contradicted those of a study in which 55% fell into the category of fair Quality of life and 20% into the category of poor quality of life, while none of the participants scored in the category of very good or excellent quality of life<sup>21</sup>. Furthermore, another study reported that 35% of participants had a good quality of life, while 46% were classified as having very good to excellent quality of life<sup>22</sup>. These studies show that quality of life measures are complicated, as individual experiences and treatment outcomes vary, reflecting the complexity of health and well-being. In the current study, regarding emotional health problems, 88.2% reported none, while 11.8% reported some. These findings were also similar to those of a study in which the majority of participants reported emotional restriction after the procedure, with a mean score of  $85.37 \pm 31.27$ , compared to the pre-mean score of  $36.75 \pm 0.78$ <sup>17</sup>. This similarity indicates that the emotional well-being of individuals across studies follows a consistent pattern.

### Comparison of Quality of Life with respect to the Sleeve gastrectomy & Lifestyle modification:

Based on the comparison of Quality-of-Life scores across the procedure and Demographic Variables, a statistically significant difference was observed in QOL scores between the two genders. In females, the mean QOL score was  $53.79 \pm 3.3$  in the sleeve gastrectomy group compared to a mean score of  $37.46 \pm 7.7$  with Lifestyle modification ( $p\text{-value} < 0.001$ ). Similarly, in males, the mean score was  $53.67 \pm 4.2$  with sleeve gastrectomy and  $40.09 \pm 9.5$  with lifestyle modification ( $p\text{-value} < 0.001$ ). These findings were consistent with the results of a previous study in which participants reported a low mean score of  $4.5 \pm 1.4$  ( $2.6 - 7.7$ ) on a scale of  $10.5 \pm 6.2$  ( $1.3 - 20.5$ )<sup>22</sup>. At the same time, the mean quality of life score without comorbidities was not significantly different, with a  $p\text{-value}$  of less than 0.05<sup>22</sup>. The mean differences in the physical and mental components among the points of both procedures were 40.6 points in the surgical procedure and 31 points in the pre-operative groups<sup>23</sup>. Likewise, another study found that the score of distal gastrectomy was +6.5 (95% CI: 0.8-12),  $p = 0.026$ , and neoadjuvant treatment scored +8.2 (95% CI: 1.8-15),  $p = 0.012$ . The minimally invasive technique scored significantly higher compared to surgical procedures<sup>24</sup>. These studies show how sleeve gastrectomy, lifestyle changes, and other surgeries impact patients' quality of life. These therapies can alter physical and mental health. Consistent differences across studies emphasize the importance of considering each treatment's specific effects, as

each intervention has benefits and drawbacks that can greatly affect a patient's quality of life.

## RECOMMENDATIONS

Future research could compare hospitals in other provinces to provide a more comprehensive understanding of quality-of-life outcomes across diverse healthcare settings. Conducting true experimental studies, such as randomized controlled trials, is recommended to generate robust evidence and identify areas for improvement in patient care. It is also recommended to organize seminars and workshops for healthcare professionals and patients to raise awareness, foster behavioral change, and improve the quality of life of individuals undergoing similar treatments.

## CONCLUSION

The mean quality of life score for sleeve gastrectomy was higher at 53.7 (SD  $\pm$  4) compared to lifestyle modification, which had a mean score of 39.45 (SD  $\pm$  9.9). A higher score reflects a better quality of life. When comparing the quality-of-life scores between the two interventions, sleeve gastrectomy and lifestyle modification, the results were statistically significant, with a p-value of less than 0.05. Additionally, there was a significant association between quality-of-life outcomes and the demographic characteristics of the participants, underscoring the influence of individual factors on the results.

**Ethical Permission:** Ziauddin University, Karachi, Pakistan, ERC approval letter No. 4960222NDNUR.

**Conflict of interest:** There is no conflict of interest between the authors.

**Financial Disclosure / Grant Approval:** No funding agency was involved in this research.

**Data Sharing Statement:** The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

## AUTHOR CONTRIBUTION

Dilshad N: Conceptualization, methodology, ethics approval, investigation (data collection), data curation, formal analysis, visualization, writing original draft, project administration.

Marshall P: Supervision, conceptual guidance, methodological oversight, validation, resources, writing review, final approval of the manuscript

Shakirullah: Co-supervision, methodology refinement, statistical guidance, interpretation of findings, writing review and editing.

## REFERENCES

1. Mramba LK, Fredericks T, Soares-da-Silva L. Investigation of non-communicable diseases prevalence, patterns, and patient outcomes in hospitalized populations: a prospective observational study in three tertiary hospitals. J

- Health Popul Nutr. 2024; 43: 128.
2. Wade AN, Jobe M, Kengne AP. Chronic noncommunicable diseases in sub-Saharan Africa: current burden and future challenges. *Lancet Glob Health*. 2024; 12(4): e512-e523. doi: 10.1016/S2214-109X(23)00549-1.
3. Lin X, Li H. Obesity: epidemiology, pathophysiology, and therapeutics. *Front Endocrinol (Lausanne)*. 2021; 12: 706978. doi: 10.3389/fendo.2021.706978
4. Rafique I, Saqib MA, Munir MA, Qureshi H, Rizwanullah KS, Khan SA et al. Prevalence of risk factors for noncommunicable diseases in adults: key findings from the Pakistan STEPS survey. *East Mediterr Health J*. 2018; 24: 33-41.
5. Kunyhamu MS, Daud A, Jusoh N. Obesity among healthcare workers: which occupations are at higher risk of being obese? *Int J Environ Res Public Health*. 2021; 18: 4381.
6. Riquelme R, Rezende LFM, Guzmán-Habinger J. Non-communicable diseases deaths attributable to high body mass index in Chile. *Sci Rep*. 2021; 11: 15500. doi: 10.1038/s41598-021-94974-z.
7. Hildebrand S, Pfeifer A. The obesity pandemic and its impact on non-communicable disease burden. *Pflugers Arch*. 2025 May; 477(5): 657-668. doi: 10.1007/s00424-025-03066-8. Epub 2025 Feb 10.
8. Reyes Matos U, Mesenburg MA, Victora CG. Socioeconomic inequalities in the prevalence of underweight, overweight, and obesity among women aged 20–49 in low- and middle-income countries. *Int J Obes (Lond)*. 2020; 44: 609-16.
9. Ibrahim S, Akram Z, Sheikh S, Zafar M, Khan N, Eraj A et al. Overweight and obesity prevalence among adults and adolescents in Gulshan-e-Iqbal, Karachi, 2018-2020. *Ann Abbasi Shaheed Hosp Karachi Med Dent Coll*. 2021; 25: 631-40.
10. Asif M, Aslam M, Altaf S, Atif S, Majid A. Prevalence and sociodemographic factors of overweight and obesity among Pakistani adults. *J Obes Metab Syndr*. 2020; 29: 58-65.
11. Hecker J, Freijer K, Hilgsmann M, Evers S. Burden of disease study of overweight and obesity: societal impact in terms of cost-of-illness and health-related quality of life. *BMC Public Health*. 2022; 22: 1-13.
12. S Safaei M, Sundararajan EA, Driss M, Boulila W, Shapi'i A. A systematic literature review on obesity: understanding the causes and consequences of obesity and reviewing machine learning approaches to predict obesity. *Comput Biol Med*. 2021; 136: 104754.
13. Reyes Matos U, Mesenburg MA, Victora CG. Socioeconomic inequalities in the prevalence of underweight, overweight, and obesity among women aged 20–49 in low- and middle-income countries. *Int J Obes (Lond)*. 2020; 44: 609-16.
14. Gulinac M, Miteva DG, Peshevska-Sekulovska M,

- Novakov IP, Antovic S, Peruhova M et al. Long-term effectiveness, outcomes and complications of bariatric surgery. *World J Clin Cases*. 2023 Jul 6; 11(19): 4504-4512. doi: 10.12998/wjcc.v11.i19.4504.
15. Kim BY, Nam H, Yoo JJ, Cho YY, Choi DH, Jung CH et al. Association between alcohol consumption status and obesity-related comorbidities in men: data from the 2016 Korean community health survey. *BMC Public Health*. 2021; 21: 1-8.
16. Seidell JC, Halberstadt J. National and local strategies in the Netherlands for obesity prevention and management in children and adolescents. *Obes Facts*. 2020; 13: 418-29.
17. Timkova V, Novak P, Horvathova K. Psychosocial distress in people with overweight and obesity: weight stigma, quality of life and mental health. *Front Psychol*. 2024; 15: 1474844. doi: 10.3389/fpsyg.2024.1474844.
18. Law S, Dong S, Zhou F, Zheng D, Wang C, Dong Z. Bariatric surgery and mental health outcomes: an umbrella review. *Front Endocrinol (Lausanne)*. 2023; 14: 1283621. doi: 10.3389/fendo.2023.1283621.
19. Castro AI, Gomez-Arbelaes D, Crujeiras AB, Granero R, Aguera Z, Jimenez-Murcia S et al. Effect of a very low-calorie ketogenic diet on food and alcohol cravings, physical and sexual activity, sleep disturbances, and quality of life in obese patients. *Nutrients*. 2018; 10: 1348.
20. Akkayaoğlu H, Çelik S. Eating attitudes, perceptions of body image and patient quality of life before and after bariatric surgery. *Appl Nurs Res*. 2020; 53: 151270. doi: 10.1016/j.apnr.2020.151270.
21. Alkassis M, Haddad FG, Gharios J, Noun R, Chakhtoura G. Quality of life before and after sleeve gastrectomy in Lebanese Popul J Obesity. 2019; 2019: 1952538. doi: 10.1155/2019/1952538.
22. Mouawad C, Nocca D, Salminen P. Quality of life after laparoscopic sleeve gastrectomy in long-term follow-up. *Obes Surg*. 2022; 32(8): 3619-3628.
23. Silva AL, Gomes LA, Pinto TS. The Impact of Bariatric Surgery on Quality of Life in Patients with Obesity: A Cross-Sectional Study. *J Clin Med*. 2022; 12(13): 4225.
24. Brenkman HJF, Tegels JJW, Ruurda JP. Factors influencing health-related quality of life after gastrectomy for cancer. *Gastric Cancer*. 2018; 21: 524-532. doi: 10.1007/s10120-017-0771-0.

