

# A Giant Well-differentiated Liposarcoma: A Case Report

Waheed Ahmed Hashmi<sup>1</sup>, Anas Hashmi<sup>2</sup>, Anam Waheed Hashmi<sup>3</sup>, Muhammad Kashif Malik<sup>4\*</sup>

## ABSTRACT

We report a rare case of an enormous well-differentiated liposarcoma (WDLPS) in a patient aged <40 years. A 39 years old Saudi male patient presented with a swelling in the upper, medial aspect of the right thigh. It was a large swelling, about 15 x 15 cm. All blood investigations were unremarkable. CT scan revealed a huge (20.1 x 20 cm) well-defined lobulated heterogeneous, hypodense subcutaneous lesion on the medial aspect of the right upper thigh, extending superiorly to the subcutaneous area of the ipsilateral natal cleft and inferiorly down to the mid-thigh level. The lesion showed multiple internal septations with a solitary spot of tiny fatty component. The entire mass was excised en bloc with its capsule. It was huge, but encapsulated. Histopathology reported a multinodular mass with infiltrative margins, measuring approximately 24 x 20 x 13 cm in greatest dimensions. The cut surface was lobulated, gelatinous, and myxoid, with grey-white, firm nodules and focal areas of haemorrhage and necrosis. Microscopic sections of the mass revealed a malignant lipomatous tumour comprising mature adipocytes, atypical spindle cells, and multivacuolated lipoblasts, embedded in a loose myxoid stroma with focal dense fibrous stroma. Few scattered mitotic figures were seen. No infiltration of the overlying skin (about 2 cm margin) was seen, and no invasion of all other margins (< 1 cm) was found. No recurrence was seen in follow-up. Documentation of LS incidence among Saudis is scarce, but it is rising in the USA and Europe. With genetic predisposition in the White population, radiation or herbicides might be responsible for this increase.

**KEYWORDS:** Liposarcomas, LS, Well-differentiated liposarcoma, WDLPS, atypical lipomatous tumor, ALT, Saudi Arabia

## INTRODUCTION

Liposarcomas (LS) are cancers derived from adipocytes or their precursor lipoblasts in soft tissues. They constitute 20% of soft tissue neoplasia<sup>1</sup>. Liposarcomas are rare mesenchymal sarcomas<sup>2</sup>. The types of liposarcomas described by WHO are atypical lipomatous tumor (ALT)/well-differentiated liposarcoma (WDLPS), dedifferentiated liposarcoma, myxoid/round cell liposarcoma (MRCLPS), and pleomorphic liposarcoma<sup>1</sup>. ALT/WDLPS can grow to enormous sizes in limbs but seldom metastasize. Limited data exist on atypical lipomatous tumor (ALT)/well-differentiated liposarcoma (WDL) in adults <40 years of age, as it is rare in this age group<sup>4</sup>. We report a case of ALT/WDLPS in a patient aged <40 years that grew to an enormous size and was managed and operated on at our institution.

## CASE DESCRIPTION

A 39 years old Saudi male patient presented in the outpatient department on December 28, 2024, with a swelling in the upper, medial aspect of the right thigh that appeared about one and a half years ago. It slowly kept increasing in size. The patient did not seek

any medical consultation for it. It was not associated with any pain or any other complaint. He had no underlying medical problem or any previous history of surgery.

On examination, a large, 15x15 cm swelling was noted on the upper medial aspect of the right thigh, with a broad base. The skin over it was intact, except for a very small area, less than 1 cm in diameter, with some skin excoriation. It was non-tender. There were no dilated veins over it. It was close to the right scrotum, but the scrotum was free and separate from the swelling. The overlying skin was not involved. It was soft, with some cystic areas, was mobile over the deeper structures/muscles, and extended posteriorly towards the perineum and anal canal. The inguinal lymph nodes were not enlarged and not palpable. It was close to the anal canal, but anal and rectal examination was normal.

All blood investigations were unremarkable. Contrast-Enhanced Computed Tomography (CECT) scan revealed a huge (20.1 x 20 cm) well-defined lobulated heterogeneous, hypodense subcutaneous lesion on the medial aspect of the right upper thigh, extending superiorly to the subcutaneous area of the ipsilateral natal cleft and inferiorly down to the mid-thigh level. The lesion showed multiple internal septations with a solitary spot of a tiny fatty component. There were no calcifications. The lesion caused displacement of adjacent thigh muscles, without definite invasion. The overlying subcutaneous tissue was intact. Adjacent bones were normal. There was no significant regional lymphadenopathy. CT scan of the chest, abdomen

<sup>1</sup>Al Iman General Hospital, Riyadh, Saudi Arabia

<sup>2</sup>General Practice Specialty Trainee, United Kingdom

<sup>3</sup>Al Faisal University, Saudi Arabia

<sup>4</sup>Prince Sultan Military Medical City, Riyadh, Saudi Arabia

Correspondence: drkmalik71@hotmail.com

doi: 10.22442/jlumhs.2026.01432

Received: 05-12-2026

Revised: 07-05-2026

Accepted: 13-05-2026

Online First: 14-07-2026



and pelvis was normal. The lungs, abdominal organs, para-aortic area, mesenteric lymph nodes, and urinary bladder were normal. A large saccular lesion, containing fluid in the medial part of the right thigh, appeared to be extending cranially.

On 30/12/2024, under spinal anesthesia, an elliptical incision was made over the mass, near its base, including ulcerated skin. It was to be excised, ensuring that the mass could be excised in toto without compromising the safety margin, and the skin could be closed later free of tension. The entire mass was excised en bloc with its capsule. It was huge but encapsulated, extending deep between the muscles of the medial compartment of the thigh, away from major nerves and vessels, and encroaching on the perineal muscles and anal sphincter without involving them. Hemostasis was secured, and the wound was closed with a drain after closing the dead space with Vicryl 3/0 sutures.

The mass was sent for histopathology. It was reported as a multinodular mass with infiltrative margins, measuring approximately 24 x 20 x 13 cm in greatest dimensions. The cut surface was lobulated, gelatinous, and myxoid, with grey-white, firm nodules and focal areas of haemorrhage and necrosis. The overlying skin ellipse measured 17 x 12 cm and was completely separate from the mass. Microscopic sections of the mass revealed a malignant lipomatous tumour comprising mature adipocytes, atypical spindle cells, and multivacuolated lipoblasts, embedded in a loose myxoid stroma with focal dense fibrous stroma. Few scattered mitotic figures were seen. No infiltration of the overlying skin (about 2 cm margin) was seen, and no invasion of all other margins (< 1 cm) was found. The diagnosis was well-differentiated liposarcoma.

The patient was referred to medical oncology for further management. He was followed up in the surgical OPD for four months. The wound initially had some serous discharge. A few clips were removed to drain the serous discharge, and the wound was treated conservatively. After drainage of the initial serous discharge, it healed well. Surgical clips were removed after 2 weeks. All investigations were repeated, including CBC, biochemistry, and LFTs, and the results were normal. He was reinvestigated by the oncologist with a total-body scan, local MRI, and PET scan, and was found to be disease-free.

Seven months later, the patient had an appointment at the Oncology Department of King Faisal Specialist Hospital in Riyadh, where they declared him disease-free.

## DISCUSSION

Liposarcomas (LS) are rare tumors. They are malignant mesenchymal sarcomas originating from soft tissues and are believed to arise from cells in the lipocyte lineage<sup>2</sup>. They constitute about 13-20% of all soft-tissue sarcomas and are the most common soft-tissue sarcomas globally<sup>2</sup>. They mostly originate in the

lower limbs, but rarely in the retroperitoneum, abdominal wall, or subcutaneous tissues<sup>1</sup>. In most cases, they appear in the lower limbs, particularly the thigh, and usually measure >11 cm<sup>8,9</sup>. Our patient also presented with this sarcoma in the thigh with similar dimensions.

In the USA, the incidence of LS is increasing, as data from 2001-2016 showed a 19% increase in LS diagnoses. In Europe, the incidence is four to five per 100,000 per year<sup>2</sup>. However, epidemiological recognition of soft tissue sarcomas in Saudi Arabia remains limited. Although they are relatively uncommon as compared to other cancers, the documentation of the incidence and characteristics of these sarcomas in the Saudi population is deficient<sup>3v</sup>. In the literature, the average age at diagnosis of LS was found to be 50 years of age<sup>2,5-9</sup>. The retroperitoneal liposarcomas usually present at a median age of 56 years; however, they can occur at any age<sup>10</sup>. Our patient, however, was quite young (39 years old).

The literature reports that men account for about 60% of cases<sup>2,5,7,8</sup>. The gender distribution of retroperitoneal sarcomas is approximately equal, although some studies have reported female predominance<sup>10</sup>. In our case, the patient was a male. In more than 80% of cases, the patients are Caucasians, less than 10% are African Americans, and the remaining belong to other races<sup>2,7</sup>. Our patient was a Saudi Arab national. One study assessed the incidence of soft-tissue sarcomas in Saudi Arabia<sup>11</sup>. The data were extracted from the Saudi Cancer Registry. This study included all the patients with primary soft-tissue sarcomas of the limbs and pelvis between January 2006 and December 2015. Out of these 659 patients, 20.5% had liposarcomas. Patients with liposarcomas had the longest mean survival (7.74 years). No widely accepted risk factors exist for liposarcomas<sup>2</sup>.

They are almost always deep-seated<sup>8</sup>. The tumor of our patient also extended deeper into the thigh and towards the perineum. It was huge and extended deep between the muscles of the medial compartment of the thigh, away from major nerves and vessels. It encroached on the perineal muscles and the anal sphincter, but did not involve them.

Treatment of WDLPS is mostly restricted to surgical resection with negative surgical margins<sup>12</sup>. They mostly do not require radiotherapy or chemotherapy. Our patient's entire tumor mass was excised en bloc with its capsule. No radiotherapy or chemotherapy was required.

On histopathological examination, these liposarcomas are usually composed of lobules of mature adipocytes, variable in size and divided into compartments by thick, fibrous bands<sup>2,8</sup>. The histopathology report of our patient showed mature adipocytes, atypical spindle cells, and multivacuolated lipoblasts embedded in a loose myxoid stroma, along with focal dense fibrous stroma, typical of WDLPS. No local or

distant metastasis was found in our patient, which is typical of this type of LS<sup>7,12</sup>.

No widely accepted risk factors for liposarcomas<sup>2</sup>. Because the etiology and risk factors are unknown, the alarming increase in liposarcoma incidence is concerning. Increased exposure to radiation or herbicides may explain this increase<sup>13</sup>. As a greater risk has been found among whites and males, there might be some genetic predisposition to liposarcoma.

## CONCLUSION

Documentation of LS incidence in the Saudi population is limited. However, its incidence has been rising in the USA and Europe. Along with genetic predisposition in the White population, radiation or herbicides might be responsible for this rise.

**Conflict of interest:** There is no conflict of interest between the authors.

**Financial Disclosure / Grant Approval:** No funding agency was involved in this research.

**Data Sharing Statement:** The corresponding author can provide the data proving the findings of this study on request. Privacy or ethical restrictions bound us from sharing the data publicly.

## AUTHOR CONTRIBUTION

Hashmi WA: Concept, revision

Hashmi A: Design, drafting

Hashmi AW: Concept, revision

Malik MK: Design, drafting

## REFERENCES

1. Finlay JR, Vijayakumar G, Braun A, Almajnooni A, Yu A, Miller I et al. Enormous myxoid/round cell liposarcoma: A case report. *Rare Tumors*. 2024 May 14; 16: 20363613241255668.
2. Jonczak E, Grossman J, Alessandrino F, Seldon Taswell C, Velez-Torres JM, Trent J. Liposarcoma: a journey into a rare tumor's epidemiology, diagnosis, pathophysiology, and limitations of current therapies. *Cancers*. 2024 Nov 18; 16(22): 3858.
3. Refai F. Soft Tissue Sarcomas: Epidemiologic Trends at King Abdulaziz University Hospital in Jeddah. *J Contemp Med Sci*. 2024 Jul 1; 10(4).
4. Waters R, Horvai A, Greipp P, John I, Demicco EG, Dickson BC et al. Atypical lipomatous tumour/ well-differentiated liposarcoma and dedifferentiated liposarcoma in patients aged ≤ 40 years: a study of 116 patients. *Histopathology*. 2019 Dec; 75(6): 833-42.
5. Lee AT, Thway K, Huang PH, Jones RL. Clinical and molecular spectrum of liposarcoma. *J Clin Oncol*. 2018 Jan 10; 36(2): 151-9.
6. Zafar R, Wheeler Y. Liposarcoma. In: *StatPearls* [Internet]. 2023 March 20. StatPearls Publishing.
7. Amer KM, Congiusta DV, Thomson JE, Elsamna S, Chaudhry I, Bozzo A et al. Epidemiology and survival of liposarcoma and its subtypes: A dual database analysis. *J Clin Orthop Trauma*. 2020 Jul 1; 11: S479-84.
8. Asano Y, Miwa S, Yamamoto N, Hayashi K, Takeuchi A, Igarashi K et al. A scoring system combining clinical, radiological, and histopathological examinations for differential diagnosis between lipoma and atypical lipomatous tumor/well-differentiated liposarcoma. *Scientific Reports*. 2022 Jan 7; 12(1): 237.
9. Zeniya T, Emori M, Tsuchie H, Teramoto A, Nagasawa H, Mizushima E. Clinicopathological and immunohistochemical analysis of the risk factors of recurrence of atypical lipomatous tumor/well-differentiated liposarcoma of the extremities. *Acta Orthopaedica Belgica*. 2024 December 1; 90: 731-7.
10. Xiao J, Liu J, Chen M, Liu W, He X. Diagnosis and prognosis of retroperitoneal liposarcoma: a single Asian center cohort of 57 cases. *J Oncology*. 2021; 2021(1): 7594027.
11. Aljuhani W, Alanazi A, Alshehri E, Alageel MK, Masuadi E, Zolaly M et al. Descriptive analysis of incidence, demographic characteristics, and survival outcomes of soft-tissue sarcoma of the extremities in Saudi Arabia. *Scientific Reports*. 2024 Nov 25; 14(1): 29123.
12. Resag A, Toffanin G, Benešová I, Müller L, Potkrajcic V, Ozaniak A et al. The immune contexture of liposarcoma and its clinical implications. *Cancers*. 2022 Sep 21; 14(19): 4578.
13. Bock S, Hoffmann DG, Jiang Y, Chen H, Il'yasova D. Increasing incidence of liposarcoma: a population-based study of national surveillance databases, 2001–2016. *Int J Environ Res Public Health*. 2020 Apr; 17(8): 2710.

